



Exposure of *Agaricus bisporus* to *Trichoderma aggressivum* f. *europaeum* leads to growth inhibition and induction of an oxidative stress response

Dejana Kosanovic ^a, Helen Grogan ^b, Kevin Kavanagh ^{a, *}

^a Department of Biology, Maynooth University, Maynooth, Co. Kildare, Ireland

^b Teagasc, Horticulture Development Department, Ashtown Research Centre, Dublin 15, Ireland

ARTICLE INFO

Article history:

Received 3 February 2020

Received in revised form

6 June 2020

Accepted 5 July 2020

Available online 23 July 2020

Keywords:

Agaricus bisporus

Trichoderma aggressivum

Proteomics

Host–pathogen interactions

ABSTRACT

Green mould disease of mushroom, *Agaricus bisporus*, is caused by *Trichoderma* species and can result in substantial crop losses. Label free proteomic analysis of changes in the abundance of *A. bisporus* proteins following exposure to *T. aggressivum* supernatant *in vitro* indicated increased abundance of proteins associated with an oxidative stress response (zinc ion binding (+6.6 fold); peroxidase activity (5.3-fold); carboxylic ester hydrolase (+2.4 fold); dipeptidase (+3.2 fold); [2Fe-2S] cluster assembly (+3.3 fold)). Proteins that decreased in relative abundance were associated with growth: structural constituent of ribosome, translation (-12 fold), deadenylation-dependent decapping of nuclear-transcribed mRNA (-3.4 fold), and small GTPase mediated signal transduction (-2.6 fold). *In vivo* analysis revealed that 10⁻⁴ *T. aggressivum* inoculum decreased the mushroom yield by 29% to 56% and 10⁻³ *T. aggressivum* inoculum decreased the mushroom yield by 68% to 100%. Proteins that increased in abundance in *A. bisporus* *in vivo* following exposure to *T. aggressivum* indicated an oxidative stress response and included proteins with pyruvate kinase activity (+2.6 fold) and hydrolase activity (+2.1 fold). The results indicate that exposure of *A. bisporus* mycelium to *T. aggressivum* *in vitro* and *in vivo* resulted in an oxidative stress response and reduction in growth.

© 2020 The Authors. Published by Elsevier Ltd on behalf of British Mycological Society. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The annual global production of edible mushrooms is estimated to be worth approximately \$42 billion dollars (Prescott et al., 2018), with the white button mushroom (*Agaricus bisporus*) being the most commonly cultivated edible mushroom in North America and Europe (Royse et al., 2017) and the main horticultural crop in the Republic of Ireland (Alexander, 2015). The most important fungal pathogens of edible mushrooms worldwide are species belonging to genera *Lecanicillium*, *Mycogone*, *Cladobotryum* and *Trichoderma* (Fletcher and Gaze, 2008; Potocnik et al., 2015). Five decades ago, *Lecanicillium fungicola*, *Mycogone perniciosa*, and *Cladobotryum* sp.,

Abbreviations: FDR, False Discovery Rates; GO, gene ontology; SSDA, statistically significant differentially abundant; DEP, differentially expressed proteins; ME, malt extract; CYM, complete yeast media; SN, supernatant; MPN, Most probably number; RH, Relative humidity; SE, Standard error.

* Corresponding author. Department of Biology, Maynooth University, Co. Kildare, Ireland.

E-mail address: Kevin.Kavanagh@mu.ie (K. Kavanagh).

<https://doi.org/10.1016/j.funbio.2020.07.003>

1878-6146/© 2020 The Authors. Published by Elsevier Ltd on behalf of British Mycological Society. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

causal agents of dry and wet bubble, and cobweb disease respectively, were the major mycopathogens damaging mushroom production (Forer et al., 1974). *Trichoderma* spp., such as *T. harzianum*, *T. atroviride*, *T. koningii*, *T. virens*, etc., have been present in mushroom industry for many years (Sinden and Hauser, 1953; Fletcher and Gaze, 2008), but in 1980s a new substrate-colonising aggressive species that severely affected mushroom production was found in Ireland (Seaby, 1987, 1996) and was subsequently named *T. aggressivum* (Samuels et al., 2002). It first appeared in two sites, in the British Isles (*T. aggressivum* f. *europaeum*) and North America (*T. aggressivum* f. *aggressivum*), and then spread worldwide causing a worldwide green mold epidemic (Rinker, 1993; Romaine et al., 1996; Geels, 1997; Hermosa et al., 1999; Mamoun et al., 2000; Hatvani et al. 2007, 2012; Szczech et al., 2008; Romero-Arenas et al., 2009; Clift and Shamshad, 2009; Kosanovic et al., 2013) and enormous economic losses (Kredics et al., 2010). This filamentous fungus is a rapidly growing, aggressive competitor that has evolved from *T. harzianum* and adapted to a specific mushroom substrate growing niche (Kredics et al., 2010; Kosanovic et al., 2015). Unlike *T. harzianum*, *T. aggressivum* colonizes the substrate for *A. bisporus*

growth and causes crop losses of between 60% and 100% (O'Brien et al., 2017; Seaby, 1996; Kredics et al., 2010). Necrotic brown spots on mushrooms may occur, while during serious outbreaks, the crop completely fails, since no fruiting bodies develop (Seaby, 1996). Disease control on mushrooms farms includes the application of only a small number of fungicides and, due to the long-term or continuous usage of fungicides, the development of fungicide-resistance has become a common phenomenon (Fletcher and Yarham, 1976; Grogan and Gaze, 2000). While *T. aggressivum* was shown to be controlled by the benzimidazole fungicide - carben-dazim in the mid-1990s (Grogan et al., 1996), resistance of *Trichoderma aggressivum* to the benzimidazole fungicide: benomyl and thiophanate-methyl was detected in North American mushroom farms in the early 2000s (Romaine et al., 2005).

The aim of this study was to characterize the interaction of *T. aggressivum* with *A. bisporus*. Microscopy and enzyme studies suggested competitive saprophytic interaction (Williams et al., 2003), but recent advances in proteomic technologies can produce novel insights into host–pathogen interactions not visible by more traditional methods. These technologies include large scale mass spectrometry-based proteomics capable of identifying and quantifying thousands of proteins across multiple samples in a single run (Zhang et al., 2013; Yates et al., 2009; Watson et al., 2015; Zhu et al., 2013; Barribeau et al., 2014). O'Brien et al., (2014) examined the proteomic response of *T. aggressivum* to *A. bisporus*, but this current study focuses on how *Agaricus* responds to the pathogen, since the pathogen *T. aggressivum* and *A. bisporus* are both present in the substrate at the same time and fungal – fungal interactions may be important factors in disease development in the crop and offer new insights into the development of novel control strategies.

2. Materials and methods

2.1. Fungal culture

T. aggressivum f. *europaeum* 100526 (acquired from the Royal Netherlands Academy of Arts and Science, Utrecht, Netherlands) preserved in liquid nitrogen culture collection of Maynooth University. PCR analysis was performed to double check the green mould re-isolated from the substrate and used as an inoculum subsequently. Three sets of primers were used (**18S INT** – 5'TAA CAA CAC GCC TGC TTA AGA'3; **TH1 INT REV** – 5'GAG AAG GCT CAG ATA GTA AAA AAT'3; **TH1 INT** – 5'CCC CCT CGC GGG GTT ATT TTT ACT'3; **EX ITS1** – 5'GTA ACA AGG TTT CCG TAG GTG'3; **EX ITS4** – 5'TTC TTT TCC TCC GCT TAT TGA TTG'3), general (ITS1/4) for *Trichoderma* spp., TH1 INT/ITS4 for *T. aggressivum*, and 18S/TH1 INT REV for subspecies *T. aggressivum* f. *europium* (Fig. S3) The results confirmed that re-isolated sample from virulence test was *T. aggressivum* f. *europaeum*. *Trichoderma* stocks were maintained on malt extract (ME) agar (Oxoid).

A. bisporus commercial strain A15 (Sylvan Inc., France) was used in this study. *A. bisporus* was maintained as spawn (sterilized rye grains colonized with *A. bisporus*) and on CYM agar plates.

2.2. Effect of *T. aggressivum* supernatants on growth of *A. bisporus* in vitro

Trichoderma cultures (10^4 conidia/ml) were grown in malt extract (ME) broth (Oxoid) at 30 °C and 200 rpm for 2 days, centrifuged and filter-sterilized through 0.2 µm syringe filter to produce supernatants. Cultures (50 ml) of *A. bisporus* (one spawn grain of A15 in CYM broth) were grown for 10 days at 25 °C and 200 rpm and supplemented with 25% v/v supernatant of *T. aggressivum* for an additional 2-, 4- or 8-d co-incubation at 25 °C

and 200 rpm. *A. bisporus* mycelium was harvested by centrifugation at 4143g for 15 min. The supernatant was discarded, and wet mass was measured.

2.3. *T. aggressivum* inoculum preparation for an in vivo test

Mushroom substrate was infected with *T. aggressivum* for an in vivo cropping trial according to the method by O'Brien et al. (2017). Briefly, inoculum was prepared for a cropping trial by inoculating Phase II mushroom substrate with *T. aggressivum* conidia. Six spawn grains were coated with $\sim 10^6$ *T. aggressivum* conidia per grain at spawning time and added to 18kg Phase II mushroom substrate, along with 90g of fresh spawn (0.5% rate). At the Teagasc Experimental Mushroom Unit (Ashtown, Co. Dublin, Ireland) mushroom growing crates (45L, 600 × 400 × 235 mm) were filled with substrate, covered and incubated under standard spawn-run conditions (25 °C and 95% RH) to produce Phase III substrate which was heavily colonised by *T. aggressivum* (Figure S1).

Samples were taken to calculate the CFU/g of *T. aggressivum* inoculum according to the method described by O'Brien et al. (2017). Homogenized substrate was diluted from 10^{-1} to 10^{-9} in sterile water and spotted onto plates. The initial fungal load in the inoculum was estimated in terms of CFUs/g of fresh weight by using the most probable number (MPN) method (Phelps, 1908). Inoculum contained 4.5×10^8 *Trichoderma* CFUs/g (Figure S2) and the *Trichoderma* present was confirmed as *T. aggressivum* f. *europaeum* by PCR (Figure S3) (O'Brien et al., 2017).

2.4. Effect of *T. aggressivum* infected substrate on growth of *A. bisporus* in vivo

T. aggressivum inoculum was added to 16 kg crates of uninfected Bulk Phase III mushroom substrate, fully colonised with *A. bisporus*, at two rates: 1 g/kg and 0.1 g/kg, equivalent to a dilution level of 10^{-3} and 10^{-4} infected substrate by weight. A control treatment of uninfected Phase III substrate was also prepared. Substrate was sampled during the first flush and the yield was harvested up to second flush. The experimental trial at the Teagasc Experimental Mushroom Unit was set up according to a randomised block design consisting of three inoculation treatments: control, 10^{-3} and 10^{-4} and five time points: day 10, 15, 20, 25 and the end of second flush (day 35) which was the end of the trial. Four replicates were prepared for each treatment combination. Compost samples for proteomics analysis were disruptively collected from the first four time point plots on days 10, 15, 20 and 25. Data on the effect of inoculum dilution level on mushroom yield for two flushes was collected from the remaining intact plots (day 35). Disease symptoms: green mould occurrence on casing and compost, deformities and spots on fruiting bodies, and pilei colorimetry assay (Chroma Meter CR-400, Konica Minolta, Sensing, INC., Japan) were also recorded from these plots.

2.5. Label free quantitative proteomics of *A. bisporus* treated with *T. aggressivum*

Proteins were extracted according to Maher et al. (2018) from *A. bisporus* mycelium which was grown in vitro for 10-d and then supplemented with *T. aggressivum* supernatant (n = 4) for an additional 4-d. *A. bisporus* proteins were also extracted from compost mycelium (in vivo testing) which was grown together with *T. aggressivum* at two different concentrations, 10^{-3} and 10^{-4} inoculum (n = 4), for 25-d in a mushroom growing room. The Bradford method was used to quantify proteins for acetone precipitation overnight. Samples were centrifuged at 14500 g for 10 min, and the pellet was re-suspended in 25 µl of 6 M urea, 2 M

thiourea and 0.1 M Tris–HCl buffer (pH 8.0). Proteins were reduced with dithiothreitol (0.5 M DTT), alkylated with iodoacetamide (0.55 M IAA) and digested with sequence grade trypsin (Promega, Ireland) at a trypsin: protein ratio of 1:40, overnight at 37 °C. Tryptic peptides were purified for mass spectrometry using C18 spin columns (Medical Supply Company, Ireland) and 0.75 µg of peptide mix was eluted onto a QExactive (ThermoFisher Scientific, USA) high resolution accurate mass spectrometer connected to a Dionex Ultimate 3000 (RSLCnano) chromatography system. Peptides were separated by an increasing acetonitrile gradient from 2% to 40% on a Biobasic C18 Picofrit column (100 mm length, 75 mm ID), using a 120-min reverse phase gradient at a low rate of 250 nl/min. A full MS scan of range 200–2000 was followed to select the 15 most intense ions prior to MS/MS. Identification of proteins from this data was performed using the Andromeda search engine in Max-Quant (version 1.2.2.5) to correlate against a *A. bisporus* database downloaded from www.uniprot.org.

The MS proteomic data and MaxQuant search output files have been added to the ProteomeXchange Consortium (Côté et al., 2012) via the PRIDE partner repository with the dataset identifier PXD016719. The Perseus software package (v. 1.5.5.3) was used for results processing, statistical analyses and graphics generation. LFQ intensities were log₂-transformed and ANOVA of significance and t-tests between the treated groups was performed using a p-value of 0.05 and significance was determined using FDR correction (Benjamini and Hochberg, 1995). Proteins which had non-existent values (suggestive of absence or very low abundance in a sample) were also used in statistical analysis. Proteins found to be absent (below the level of detection) in one or more treatments and present (above the level of detection) in three or fewer treatments were also used in statistical analysis of the total differentially expressed group following imputation of the zero values with values that simulate low abundant proteins.

These proteins were subjected to Gene Ontology (GO) analysis by Blast2GO software tool (<https://www.blast2go.com/>). Statistically enriched GO descriptors were examined to identify clusters of proteins enriched within statistically significant differentially abundant (SSDA) protein lists ($P < 0.05$).

2.6. Statistical analysis

Experiments performed *in vitro* were carried out on three independent occasions, in four replicates and results are expressed as the mean ± SE. Proteomic and yield data from *in vivo* testing had four replicates. Significant differences were considered at $P < 0.05$.

3. Results

3.1. Assessment of effect of *T. aggressivum* supernatants on *A. bisporus* growth *in vitro*

The effect of *T. aggressivum* supernatants on *A. bisporus* growth *in vitro* was assessed after 10-day incubation of *A. bisporus* at 25 °C and additional 2-, 4- and 8-d of co-incubation with 25% v/v *T. aggressivum* filter-sterilized 48h supernatant or MEB (control). After two days, *T. aggressivum* supernatant inhibited growth by 27.9% where average growth for the control was 1.64 ± 0.22 g and for the treatment was 1.18 ± 0.11 g (Fig. 1). After four days, *T. aggressivum* supernatant statistically significantly inhibited growth by 48% ($P < 0.05$, $t = 3.807$, $df = 6$), where average growth for the control was 2.12 ± 0.22 g and for the treatment was 1.11 ± 0.15 g (Fig. 1). After eight days *T. aggressivum* supernatant inhibited growth by 13.16%, average growth for the control was 4.10 ± 0.34 g and for the treatment was 3.56 ± 0.29 g (Fig. 1). Only




Fig. 1. *A. bisporus* growth [g] in CYM after 12-, 14- and 18-d incubation at 25 °C, containing 25% v/v of either: MEB (Control) or *T. aggressivum* 48h supernatant (+*T. aggressivum* SN) for the last 2-, 4- and 8-d of incubation (+2, +4 + 8). 12-d: $P = 0.116$, $t = 1.836$, $df = 6$; 14-d: $**P = 0.01$, $t = 3.807$, $df = 6$; 18-d: $P = 0.258$, $t = 1.217$, $df = 8$.

four-day co-incubation was found to significantly ($P = 0.05$) inhibit *A. bisporus* growth (Fig. 1).

3.2. Proteomic analysis of *A. bisporus* response after exposure to *T. aggressivum* supernatant *in vitro*

Proteomic analysis was performed on one set of samples, where growth of *A. bisporus* has been significantly affected. Proteins were extracted from four replicate samples of *A. bisporus* exposed to *T. aggressivum* supernatant for 4 days and prepared for label free quantitative proteomics. In total 17654 peptides were identified, representing 1226 proteins. According to Perseus analysis a total of 73 *A. bisporus* proteins in the *T. aggressivum* +4 day treatment were determined to be differentially abundant (analysis of variance [ANOVA], $P < 0.05$), 36 increased and 37 decreased in relative abundance with a fold change value of >1.5 . A total of 280 *A. bisporus* proteins were deemed exclusive (i.e., with LFQ intensities present in all four replicates of treatment and absent in all four replicates of control). These proteins were also used in statistical analysis of the total levels corresponding to the differentially expressed group members following imputation of the zero values as described above. After data imputation, these proteins were included in the subsequent statistical analysis.

A principal-component analysis (PCA) performed on all filtered proteins distinguished the control and *T. aggressivum* -treated samples (Figure S4). Label free proteomic analysis of changes in the relative abundance of *A. bisporus* proteins, compared to the control, following exposure of cells to *T. aggressivum* supernatant was performed, and results indicated that exposure lead to an oxidative stress response (proteins that increased in relative abundance: zinc ion binding (+6.6 fold); peroxidase activity (+5.3 fold); carboxylic ester hydrolase (+2.4 fold); biosynthetic process (+3.1 fold); 60S ribosomal protein L36 (+3.3 fold); dipeptidase (+3.2 fold) [2Fe-2S] cluster assembly (+3.3 fold)) (Fig. 2).

The proteins that were decreased in relative abundance in *A. bisporus* exposed to *T. aggressivum* supernatant were associated with growth: structural constituent of ribosome, translation (-12 fold), deadenylation-dependent decapping of nuclear-transcribed mRNA (-3.4 fold), superoxide dismutase (-3 fold), metal ion binding (-2.7 fold), small GTPase mediated signal transduction (-2.6 fold), deoxyribonucleotide catabolic process (-2.6 fold), GTP binding (-2.7 fold), glycine cleavage system P protein (-2.3 fold), and proteasome subunit beta (-2.3 fold) (Fig. 2).

Proteins were subjected to Gene Ontology (GO) analysis by Blast2GO software tool. Several GO terms belonging to biological process (oxidation-reduction response to stress, metabolic and



Fig. 4. Mushroom productivity during the first flush. A - Control; B - 10^{-4} inoculation rate; C - 10^{-3} inoculation rate. Note the presence of green mould, diseased mushrooms and reduced cropping area in 10^{-4} and 10^{-3} inoculation treatments compared to the control.

samples obtained on days 20 and 25. The results obtained on day 25 indicated that a total of 35 peptides were identified, representing 10 proteins. According to Perseus analysis a total of 7 *A. bisporus* proteins in *T. aggressivum* 10^{-4} infection level compost were determined to be differentially abundant (analysis of variance [ANOVA], $P < 0.05$), 1 increased and 6 decreased in relative abundance with a fold change value of >1.5 . One *A. bisporus* protein was deemed exclusive (i.e., with LFQ intensities present in all four replicates of treatment and absent in all four replicates of control) and was used in statistical analysis of the total levels corresponding to the differentially expressed group members following imputation of the zero values as described above. After data imputation, this protein was included in the subsequent statistical analysis, and results indicated that exposure lead to decrease in relative abundance of proteins involved in hydrolase activity (+4.3 fold) (Figure S8).

Label free proteomic analysis of changes in the abundance of *A. bisporus* proteins at day 25, following exposure of cells to *T. aggressivum* 10^{-3} inoculum was performed, and in total 41 peptides were identified, representing 10 proteins. According to Perseus analysis a total of 5 *A. bisporus* proteins in this *T. aggressivum* treatment were determined to be differentially abundant (analysis of variance [ANOVA], $P < 0.05$), 4 increased and 1 decreased in relative abundance with a fold change value of >1.5 . Results indicated that exposure lead to an oxidative stress response (proteins that increased in relative abundance: kinase activity (+2.6 fold), magnesium ion binding +2.6 fold), potassium ion binding, pyruvate kinase activity (+2.6 fold); carbohydrate metabolic process and beta-galactosidase activity (+2.2 fold); hydrolase activity [GO:0016787] (+2.1 fold); and hydrolase activity [GO:0016787] (+1.8 fold)) (Figure S9). The protein that decreased in relative abundance in *A. bisporus* treated with *T. aggressivum* 10^{-3} inoculum was involved in polysaccharide catabolic process, and the lyase activity (-3 fold) (Figure S9).

Proteins were subjected to Gene Ontology (GO) analysis by the Blast2GO software tool. Several GO terms belonging to molecular function (hydrolase activity), cellular components (intrinsic component of membrane) were significantly enriched within the datasets in both levels of infection. Hydrolases and oxidoreductases were the most enriched enzymes in 10^{-4} inoculated plots whereas in 10^{-3} infected plots hydrolases and transferases were most

abundant amongst proteins identified in *A. bisporus* treated with *T. aggressivum* (Figure S10 a, b, c).

4. Discussion

The aim of the work presented here was to assess the response of *A. bisporus* to exposure to *T. aggressivum* both *in vitro* and *in vivo* in order to gain a better understanding of how they interact and why *A. bisporus* growth is so inhibited. The reduction in biomass caused by exposure of *A. bisporus* to *T. aggressivum* supernatant in the laboratory correlated with the yield decrease in mushroom growing facility following the green mould infection. *In vitro* results showed that *T. aggressivum* supernatants reduced *A. bisporus* growth at all time points, but the statistically significant ($P = 0.05$) decrease of 48% was found after four days of co-incubation. The cause of the reduced effect of mould supernatant may be due to the absence of a live organism in the supernatant and *A. bisporus* ability to overcome the impact of the one-time addition of supernatant. The transient effect of supernatant is mirrored in the proteomics analysis performed on 2- and 8-day (Fig S11 a,b) samples which showed minor changes vs control. However, proteomic response of mycelia after 4-day co-incubation with *T. aggressivum* supernatant (Fig. 2) exhibited increases in proteins associated with oxidative stress response and distinguishes from the control. *In vivo* experiments, with continuous exposure to *T. aggressivum*, demonstrated that mushroom yield is vulnerable to both levels of *T. aggressivum* infection and the extent of crop loss directly correlates with the green mould infection level. A higher dose of *T. aggressivum* caused more severe disease symptoms in the substrate, on the crop surface and on fruiting bodies and resulted in lower yield, with reductions of 37% at 10^{-4} level and 76% at 10^{-3} level of infection compared to the controls.

Proteomic assessment revealed that both groups of samples, laboratory harvested wet mass (*in vitro* experiments) and mushroom growing compost samples (*in vivo* experiments), showed *A. bisporus* increased abundance of oxidative stress proteins (peroxidase, hydrolase, kinase and ion binding activity) when cultured in the presence of *T. aggressivum*. Also, proteins associated with growth were found to be reduced in abundance (polysaccharide catabolic process, structural constituent of ribosome, translation, deadenylation-dependent de-capping of nuclear-transcribed mRNA, small GTPase mediated signal transduction,

deoxyribonucleotide catabolic process, GTP binding, glycine cleavage system P protein and proteasome subunit beta), indicating that the two fungal organisms not only compete for space and nutrients (Williams et al., 2003) but do interact more as pathogen and host.

T. aggressivum does produce secondary metabolites, and volatile organic compounds found to negatively affect mushroom yield (Kredics et al., 2010). However, *A. bisporus* extracts have been found to promote *T. aggressivum* growth and conidia germination which was not seen in other species of genus *Trichoderma* (Mamoun et al., 2000; Mumpuni et al., 1998). Diverse opinions about the *T. aggressivum* f. *europaeum* development in compost could be found. Thus, *T. aggressivum* mycelium is barely visible in compost without the presence of *A. bisporus*, suggesting that aggressive *Trichoderma* needs the presence of *A. bisporus* to stimulate growth (Seaby et al., 1996; Rinker, 1996; Romaine et al., 1996). Mamoun et al. (2000), discovered that the mycelium of *A. bisporus* is required for the induction of intensive sporulation of *T. aggressivum* f. *europaeum*.

The *Trichoderma* genus stands out among other Ascomycotina members for its abilities to synthesize numerous secondary metabolites, *in vivo* and *in vitro*, and many of its enzymes are used in industrial and biotechnology processes (Sivasithamparam and Ghisalbert, 1998; Kubicek et al., 2009). In accordance with EU regulations for environmental protection and health, the Sustainable Use of Pesticides Directive (SUD) 2009/128/EC regulates the use of pesticides in agriculture and advocates an integrated pest management (IPM) approach to pesticide use whereby non-chemical biocontrol approaches to pest and disease control should be used where possible. Some *Trichoderma* spp. with antagonistic antifungal properties are used as biocontrol agents in plant protection since early 1930s (Weindling, 1932; Harman et al., 2010; Lorito et al., 2010). Among the first such antifungal biocontrol substances isolated from certain *Trichoderma* spp. was 6-pentyl- α -pyrone, compound that has been synthetically made after it was first isolated from *T. viride* (Collins and Hilim, 1972). The isolated volatile compound has a characteristic coconut smell and is present in *T. viride*, *T. atroviride* and *T. koningii* (Bisby, 1939; Dodd et al., 2003). While the antifungal properties of *Trichoderma* spp may be beneficially used for the control of fungal pathogens of plant-based crops, they are not desirable in mushroom crop production, where the crop is itself a fungus. Other mushroom crops such as *Pleurotus ostreatus* and *Lentinula edodes* also suffer from *Trichoderma* infections, highlighting the aggressive antifungal nature of this genus (Qiu et al., 2017; Wang et al., 2016).

Previous studies have suggested that *T. aggressivum* is a competitor for space and is a saprophytic contaminant in compost for mushroom cultivation rather than mycoparasite (Williams et al., 2003; O'Brien et al., 2014). Even though *T. aggressivum* produces stress tolerance and longevity proteins when in contact with *A. bisporus* and Phase 3 mushroom substrate (O'Brien et al. 2014, 2017; Williams et al., 2003; Hatvani et al., 2010) it should be considered as a true mycopathogen as the results presented here show a direct effect of *T. aggressivum* supernatant and hyphae on *A. bisporus* development and the induction of an oxidative stress response. This understanding is an important epidemiological insight further unmasking the *T. aggressivum* diseases processes which may enable development of novel ways of prevention, control and treatment of this serious disease of the mushroom crop.

Compliance with ethical standards

This article does not contain any study with human participants or animals performed by the authors.

Declaration of competing interest

The authors have no conflicts of interest to declare.

Acknowledgements

DK is a Postdoctoral Fellow supported by Irish Research Council, GOIPD/2018/115. Q-Exact mass spectrometer was funded under the SFI Research Infrastructure Call 2012; Grant Number: 12/RI/2346. Marilyn Bonham, Department of Plant Pathology and Environmental Microbiology, Pennsylvania State University, PA, USA for helping with the crop work, setting of the experiment and gathering data. Irene Marongiu, Teagasc, Horticulture Development Department, Ashtown Research Centre, for setting and monitoring the crop in the Mushroom Unit and collecting data. Brian McGuinness, Teagasc Plant Pathology Lab for helping with PCR analysis. Eoin O'Connor, Maynooth University Bioinformatics Lab, for helping with proteomic results analysis and virulence test setting. Justina Jablokov, Maynooth University, for help with protein extractions.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.funbio.2020.07.003>.

References

- Alexander, S., 2015. Horticulture in Ireland. Teagasc Res 10 (3), 18–19. <https://www.teagasc.ie/media/website/publications/2015/TResearch-Autumn-15.pdf>.
- Barribeau, S.M., Sadd, B.M., DuPlessis, L., Schmid-Hempel, P., 2014. Gene expression differences underlying genotype-by-genotype specificity in a host–parasite system. Proc. Natl. Acad. Sci. Unit. States Am. 111, 3496–3501.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J. Roy. Stat. Soc. B 57 (1), 289–300.
- Bisby, G.R., 1939. *Trichoderma viride* pers. Ex fries, and notes on hypocrea. Trans. Br. Mycol. Soc. 23, 149–168.
- Collins, R.P., Halim, A.F., 1972. Characterisation of the major aroma constituent of the fungus *Trichoderma viride* (Pers.). J. Agric. Food Chem. 20, 437–438.
- Côté, R.G., Griss, J., Dienes, J.A., Wang, R., Wright, J.C., van den Toorn, H.W.P., et al., 2012. The PRoteomics IDentification (PRIDE) converter 2 framework: an improved suite of tools to facilitate data submission to the PRIDE database and the ProteomeXchange consortium. Mol. Cell. Proteomics 11, 1682–1689.
- Clift, A.D., Shamshad, A., 2009. Modeling mites, moulds and mushroom yields in the Australian mushroom industry. In: Anderssen, R.S., Braddock, R.D., Newham, L.T.H. (Eds.), Proceedings of the 18th World IMACS/MODSIM09 Congress Cairns: IMACS/MODSIM, pp. 491–497.
- Dodd, S.L., Lieckfeldt, E., Samuels, G.J., 2003. *Hypocrea atroviridis* sp. nov., the tel-eomorph of *Trichoderma atroviride*. Mycologia 95 (1), 27–40.
- Fletcher, J.T., Gaze, R.H., 2008. Mushroom Pest and Disease Control. Manson Publishing, London.
- Fletcher, J.T., Yarham, D.J., 1976. The incidence of benomyl tolerance in *Verticillium fungicola*, *Mycogone perniciosa* and *Hypomyces rosellus* in mushroom crops. Ann. Appl. Biol. 84, 343–353.
- Forer, L.B., Wuest, P.J., Wagner, U.R., 1974. Occurrence and economic impact of fungal diseases of mushrooms in Pennsylvania. Plant Dis. Rep. 54, 987–991.
- Geels, F.P., 1997. Rondetafel – bijeenkomst over *Trichoderma*. Champignoncultuur 41, 13.
- Grogan, H.M., Gaze, R.H., 2000. Fungicide resistance among *Cladobotryum* spp.: causal agent of cobweb disease of the edible mushroom *Agaricus bisporus*. Mycol Res 104, 357–364.
- Grogan, H.M., Noble, R., Gaze, R.H., Fletcher, J.T., 1996. Control of *Trichoderma harzianum*: a weed mould of mushroom cultivation. In: Proceedings of Brighton Crop Protection Conference—Pests and Disease. BCPC, Farnham, Surrey, UK, pp. 337–342.
- Harman, G.E., Obregón, M.A., Samuels, G., Lorito, M., 2010. Changing models of biocontrol in the developing and developed world. Plant Dis. 94 (8), 928–939.
- Hatvani, L., García Jimenez, L., Naeimi, S., Czifra, D., Urbán, P., Manczinger, L., Vágvolgyi, C., Kredics, L., 2010. A challenge to mushroom growers: the green mould disease of cultivated champignons. In: Méndez-Vilas, A. (Ed.), Current Research, Technology and Education, Topics in Applied Microbiology and Microbial Biotechnology. Publisher, FORMATEX, pp. 295–305.
- Hermosa, M.R., Grondona, I., Monte, E., 1999. Isolation of *Trichoderma harzianum* Th2 from commercial mushroom compost in Spain. Plant Dis. 83, 591.

- Kosanovic, D., Potocnik, I., Duduk, B., Vukojevic, J., Stajic, M., Rekanovic, E., Milijasevic-Marcic, S., 2013. *Trichoderma* species on *Agaricus bisporus* farms in Serbia and their biocontrol. *Ann. Appl. Biol.* 163, 218–230.
- Kosanovic, D., Potocnik, I., Vukojevic, J., Stajic, M., Rekanovic, E., Stepanovic, M., Todorovic, B., 2015. Fungicide sensitivity of *Trichoderma* spp. from *Agaricus bisporus* farms in Serbia. *J. Environ. Sci. Health Part B Pestic. Food Contam. Agric. Wastes* 50 (8), 607–613.
- Kredics, L., Jimenez, L.G., Naeimi, S., Czifra, D., Urban, P., Manczinger, L., Vagvolgyi, C., Hatvani, L., 2010. A challenge to mushroom growers: the green mould disease of cultivated champignons. *Current Res Technol and Education Topics in Appl Microbiology and Microbial Biotechnology* 295–305.
- Kubicek, C.P., Mikus, M., Schuster, A., Schmolli, M., Seiboth, B., 2009. Metabolic engineering strategies for the improvement of cellulase production by *Hypocrea jecorina*. *Biotechnol. Biofuels* 1, 2–19.
- Lorito, M., Woo, S.L., Harman, G.E., Monte, E., 2010. Translational research on *Trichoderma*: from 'omics to the field. *Annu. Rev. Phytopathol.* 48, 395–417. <https://doi.org/10.1146/annurev-phyto-073009-114314>.
- Maier, A., Staunton, K., Kavanagh, K., 2018. Analysis of the effect of temperature on protein abundance in *Demodex*-associated *Bacillus oleronius*. *Pathogens and Dis.* <https://doi.org/10.1093/femspd/fty032>.
- Mamoun, M.L., Iapicco, R., Savoie, J.M., Olivier, J.M., 2000. Green mould disease in France: *Trichoderma harzianum* Th2 and other species causing damage on mushroom farms. *Mushroom Sci.* 15, 625–632.
- Mumpuni, A., Sharma, H.S.S., Brown, A., 1998. Effect of metabolites produced by *Trichoderma harzianum* biotypes and *Agaricus bisporus* on their respective growth radii in culture. *Appl. Environ. Microbiol.* 64 (12), 5053–5056.
- O'Brien, M., Grogan, H., Kavanagh, K., 2014. Proteomic response of *Trichoderma aggressivum* to *Agaricus bisporus* tissue and mushroom compost. *Fungal Biol* 118, 785–791.
- O'Brien, M., Kavanagh, K., Grogan, H., 2017. Detection of *Trichoderma aggressivum* in bulk phase III substrate and the effect of *T. aggressivum* inoculum, supplementation and substrate mixing on *Agaricus bisporus* yields. *Eur. J. Plant Pathol.* 147 (1), 199–209.
- Phelps, E.B., 1908. A method for calculating the number of *B. coli* from the results of dilution tests. *Am. J. Hyg.* 4, 141–145.
- Potocnik, I., Stepanovic, M., Rekanovic, E., Todorovic, B., Milijasevic-Marcic, S., 2015. Disease control by chemical and biological fungicides in cultivated mushrooms: button mushroom, oyster mushroom and shiitake. *Pesticides and Phytomed* 30 (4), 201–208.
- Prescott, T.J., Wong, B., Panaretou, E., Boa, A., Bond, et al., 2018. Useful fungi. In: Willis, K. (Ed.), *State of the World's Fungi*, Report. Royal Botanic Gardens, Kew., pp. 24–31.
- Qiu, Z., Wu, X., Zhang, J., Huang, C., 2017. High temperature enhances the ability of *Trichoderma asperellum* to infect *Pleurotus ostreatus* mycelia. *PLoS One* 12 (10), e0187055. <https://doi.org/10.1371/journal.pone.0187055>.
- Rinker, D.L., 1993. Disease management strategies for *Trichoderma* mould. *Mushroom World* 4, 3–5.
- Rinker, D.L., 1996. *Trichoderma* disease: progress toward solutions. *Mushroom World* 7, 46–53.
- Romaine, C.P., Royse, D.J., Wuest, P.J., Beyer, D.M., 1996. Mushroom green mould: cause, edaphic factors and control. *Mushroom News* 44, 20–23.
- Romaine, C.P., Royse, D.J., Schlaghauser, C., 2005. Superpathogenic *Trichoderma* resistant to topsin M found in Pennsylvania and Delaware. *Mushroom News* 53, 6–9.
- Romero-Arenas, O., Lara, M.H., Huato, M.A.D., Hernandez, F.D., Victoria, D.A.A., 2009. The characteristics of *Trichoderma harzianum* as a limiting agent in edible mushrooms. *Rev. Colomb. Biotechnol.* 11, 143–151.
- Royse, D.J., Baars, J., Tan, Q., 2017. Current overview of mushroom production in the world. In: Zied, D.C., Pardo-Giménez, A. (Eds.), *Edible and Medicinal Mushrooms*, first ed. Technology and applications, pp. 2–13.
- Samuels, G.J., Dodd, S.L., Gams, W., Castlebury, L.A., Petrini, O., 2002. *Trichoderma* species associated with the green mold epidemic of commercially grown *Agaricus bisporus*. *Mycologia* 94, 147–170.
- Sivasithamparam, K., Ghisalberti, E.L., 1998. Secondary metabolism in *Trichoderma* and *gliocladium*. In: Kubicek, C.P., Harman, G.E. (Eds.), *Trichoderma and Gliocladium*. Vol. 1. Basic Biology, Taxonomy and Genetics. Taylor and Francis Ltd., London, pp. 139–191.
- Szczecz, M., Staniaszek, M., Habdas, H., Uliński, Z., Szymanski, J., 2008. *Trichoderma* spp.—The cause of green mould on polish mushroom farms. *Veg. Crops Res. Bull.* 69, 105–114.
- Seaby, D.A., 1987. Infection of mushroom compost by *Trichoderma* species. *Mushroom J.* 179, 355–361.
- Seaby, D.A., 1996. Investigation of the epidemiology of green mould of mushroom (*Agaricus bisporus*) compost caused by *Trichoderma harzianum*. *Plant Pathol.* 45, 913–923.
- Sinden, J., Hauser, E., 1953. Nature and control of three mildew diseases of mushrooms in America. *Mushroom Sci.* 2, 177–180.
- Wang, G., Cao, X., Ma, X., Guo, M., Liu, C., Yan, L., Bian, Y., 2016. Diversity and effect of *Trichoderma* spp. associated with green mold disease on *Lentinula edodes* in China. *Microbiologyopen* 5 (4), 709–718. <https://doi.org/10.1002/mbo3.364>.
- Watson, A.K., Williams, T.A., Williams, B.A., Moore, K.A., Hirt, R.P., Embley, T.M., 2015. Transcriptomic profiling of host-parasite interactions in the microsporidian *Trachipleistophora hominis*. *BMC Genom.* 16, 975–983.
- Weindling, R., 1932. *Trichoderma lignorum* as a parasite of other soil fungi. *Phytopathology* 22, 837–845.
- Williams, J., Clarkson, J.M., Mills, P.R., Cooper, R.M., 2003. Saprotrophic and myco-parasitic components of aggressiveness of *Trichoderma harzianum* groups toward the commercial mushroom *Agaricus bisporus*. *Appl. Environ. Microbiol.* 69 (7), 4192–4199. <https://doi.org/10.1128/AEM.69.7.4192-4199.2003>.
- Yates, J.R., Ruse, C.I., Nakorchevsky, A., 2009. Proteomics by mass spectrometry: approaches, advances, and applications. *Annu. Rev. Biomed. Eng.* 11, 49–79.
- Zhang, Y., Fonslow, B.R., Shan, B., Baek, M.-C., Yates III, J.R., 2013. Protein analysis by shotgun/bottom-up proteomics. *Chem. Rev.* 113, 2343–2394.
- Zhu, J.-Y., Yang, P., Zhang, Z., Wu, G.-X., Yang, B., 2013. Transcriptomic immune response of *Tenebrio molitor* pupae to parasitization by *Scleroderma guani*. *PLoS One* 8, e54411.