



In-depth quantitative profiling of post-translational modifications of Timothy grass pollen allergome in relation to environmental oxidative stress

Katarina Smiljanic^a, Ivana Prodic^b, Danijela Apostolovic^c, Anka Cvetkovic^d, Djordje Veljovic^e, Jelena Mutic^{a,f}, Marianne van Hage^c, Lidija Burazer^g, Tanja Cirkovic Velickovic^{a,f,h,i,*}

^a University of Belgrade-Faculty of Chemistry, Centre of Excellence for Molecular Food Sciences and Department of Biochemistry, Belgrade, Serbia

^b Innovation Center Ltd, University of Belgrade-Faculty of Chemistry, Belgrade, Serbia

^c Karolinska Institute, Department of Medicine, Solna, Stockholm, Sweden

^d Institute of Public Health of Belgrade, Belgrade, Serbia

^e University of Belgrade-Faculty of Technology and Metallurgy, Belgrade, Serbia

^f Ghent University Global Campus, Incheon, South Korea

^g Institute of Immunology, Virology and Sera Production, Torlak Institut, Belgrade, Serbia

^h Ghent University, Faculty of Bioscience Engineering, Ghent, Belgium

ⁱ Serbian Academy of Sciences and Arts, Belgrade, Serbia

ARTICLE INFO

Handling Editor: Martí Nadal

Keywords:

Timothy grass pollen

Label free relative quantification

Oxidative post-translational modification

Air-related traffic pollution

Allergy

Heavy metal pollution

ABSTRACT

An association between pollution (e.g., from traffic emissions) and the increased prevalence of respiratory allergies has been observed. Field-realistic exposure studies provide the most relevant assessment of the effects of the intensity and diversity of urban and industrial contamination on pollen structure and allergenicity. The significance of in-depth post-translational modification (PTM) studies of pollen proteomes, when compared with studies on other aspects of pollution and altered pollen allergenicity, has not yet been determined; hence, little progress has been made within this field. We undertook a comprehensive comparative analysis of multiple polluted and environmentally preserved *Phleum pratense* (Timothy grass) pollen samples using scanning electron microscopy, in-depth PTM profiling, determination of organic and inorganic pollutants, analysis of the release of sub-pollen particles and phenols/proteins, and analysis of proteome expression using high resolution tandem mass spectrometry. In addition, we used quantitative enzyme-linked immunosorbent assays (ELISA) and immunoglobulin E (IgE) immunoblotting. An increased phenolic content and release of sub-pollen particles was found in pollen samples from the polluted area, including a significantly higher content of mercury, cadmium, and manganese, with irregular long spines on pollen grain surface structures. Antioxidative defense-related enzymes were significantly upregulated and seven oxidative PTMs were significantly increased (methionine, histidine, lysine, and proline oxidation; tyrosine glycosylation, lysine 4-hydroxy-2-nonenal adduct, and lysine carbamylation) in pollen exposed to the chemical plant and road traffic pollution sources. Oxidative modifications affected several Timothy pollen allergens; Phl p 6, in particular, exhibited several different oxidative modifications. The expression of Phl p 6, 12, and 13 allergens were downregulated in polluted pollen, and IgE binding to pollen extract was substantially lower in the 18 patients studied, as measured by quantitative ELISA. Quantitative, unrestricted, and detailed PTM searches using an enrichment-free approach pointed to modification of Timothy pollen allergens and suggested that heavy metals are primarily responsible for oxidative stress effects observed in pollen proteins.

Abbreviations: 1D, one dimensional; 2D, two dimensional; BSA, bovine serum albumin; Cys, cysteine amino acid; ELISA, enzyme-linked immunosorbent assay; EPA, Environmental Protection Agency; GAE, gallic acid equivalents; His, histidine amino acid; HNE, 4-hydroxy-2-nonenal (Michael adduct); IgE, immunoglobulin E; ICP-MS, inductively coupled plasma mass spectrometry; ICP-OES, inductively coupled plasma optical emission spectrometry; kDa, kilodaltons; LFQ, label free quantification; Lys, lysine amino acid; Met, methionine amino acid; MS/MS, tandem mass spectrometry; nLC-MS/MS, nano-liquid chromatography coupled to tandem mass spectrometry; P1, location of environmentally preserved Timothy grass pollen; P2, location of air-related traffic and factory polluted Timothy grass pollen; PAHs, polycyclic aromatic hydrocarbons; Phl p, code for allergens from *Phleum pratense* pollen; PTM, post-translational modification; SDS-PAGE, sodium dodecyl sulphate polyacrylamide gel electrophoresis; SEM, scanning electron microscopy; SPPs, sub-pollen particles; ROS, reactive oxygen species; RT, room temperature (between 20 °C and 25 °C); tPBS, Tween 20 phosphate buffered saline; Tyr, tyrosine amino acid; XIC, extracted ion chromatogram

* Corresponding author at: University of Belgrade-Faculty of Chemistry, Center of Excellence for Molecular Food Sciences and Department of Biochemistry, Studentski trg 12-16, 11 000 Belgrade, Serbia.

E-mail addresses: katarinas@chem.bg.ac.rs (K. Smiljanic), tcirkov@chem.bg.ac.rs, Tanja.Velickovic@ghent.ac.kr (T. Cirkovic Velickovic).

<https://doi.org/10.1016/j.envint.2019.03.001>

Received 4 October 2018; Received in revised form 2 February 2019; Accepted 1 March 2019

Available online 08 March 2019

0160-4120/ © 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In order to explain observed differences in allergy prevalence rates, epidemiological studies have demonstrated an association between pollution, such as traffic emissions, and an increased prevalence of allergic symptoms. In addition to explaining the observed differences in allergy prevalence between rural, environmentally preserved areas and urban areas, epidemiological studies have also shown an association between pollution (e.g., traffic emissions) and an increase in the prevalence of respiratory allergies (Ishizaki et al., 1987; Kramer et al., 2000; Parker et al., 2009; Ruokolainen et al., 2015). However, the direct effects of air and soil pollution on the structure and physiology of the allergen carrier, i.e. pollen, including the mechanisms that lead to the altered allergenicity and the potential for enhanced symptom severity in patients suffering from pollen allergy, still warrant further research. This is because the complexity of the interactions between pollen species, contamination types, and exposure duration, as well as the methodology used to assess pollution effects all influencing the results obtained (Senechal et al., 2015). The contributions of studies that have undertaken artificial, experimentally controlled approaches to determine the impact of air and heavy metal pollution on pollen structure and/or allergenicity are important (Senechal et al., 2015). However, the intensity and diversity of the effects of urban and industrial contamination on pollen has so far been the most optimally assessed and confirmed by field-realistic exposure studies (Ishizaki et al., 1987; Kramer et al., 2000; Parker et al., 2009).

Phleum pratense pollen is one of the most frequently pollen grains found in the bioaerosol masses in Europe and the USA (Motta et al., 2006; Schmidt et al., 2010). It is a widespread grass species that has high cross-reactivity and can be readily characterized with different proteomic preparations, including two dimensional (2D) immunoblot allergomes (Schmidt et al., 2010). Motta et al. (2006) exposed commercial *Phleum pratense* pollen grains to different concentrations of traffic-related air gaseous pollutants (O_3 and NO_2) separately. They found a significant increase in the proportion of grains that released sub-pollen particles (SPPs) by osmolysis, and increased the extent of severely damaged pollen (Motta et al., 2006). In a later study, the authors found that the exposure of pollen to gaseous pollutants (O_3 , NO_2 , and SO_2) individually and in combination induced a decrease in allergen immunoglobulin E (IgE) binding. They attributed this to post-translational modifications (PTMs), since the 2D gel profiles demonstrated unchanged allergenic content and abundance. The authors concluded that these results warranted further investigation (Rogerieux

et al., 2007).

Since then, in-depth PTM searches of pollen proteomes have made the least progress within this field, compared with studies that examine other aspects related to pollen pollution (e.g., germination, viability, releasing properties of SPPs, proteome expression profiles, or profiling of organic and inorganic pollutants). A review from 2015 examined > 250 studies, addressing the effects of air-borne pollutants on different pollen species and their allergenicity, mainly by in vitro experimentally controlled methods. None of these studies considered the in-depth pollen proteome PTMs, not even the studies that considered differential transcriptomes and allergenicity (El Kelish et al., 2014; Senechal et al., 2015; Zhao et al., 2017; Zhao et al., 2016). The significance of, and requirement for, in-depth (wide and deep) PTM searches has not been adequately recognized by the scientific community. One exception to this, albeit in the direction of deep PTM knowledge, is a study on the artificial nitration of Bet v 1a. In this study, tyrosine (Tyr 83) was nitrated within the calyx cavity, and this induced a shift towards the Th_2 immuno-response. This stressed the importance of protein PTMs in understanding both the molecular basis of allergenicity and connections between the environment and health at the molecular level (Ackaert et al., 2014). Therefore, the exploration and quantification of different PTMs provides the basis for novel, mechanistic insights.

Pollen pollution by heavy metals (mercury, lead, and cadmium), transition metals (nickel, chromium, and manganese), and polycyclic aromatic hydrocarbons (PAHs) has been more frequently studied from the perspective of environmental pollution and food safety, compared with pollen allergenicity and the severity of allergy symptoms. The very rare exception to this is a study by Aina et al. (2010), who showed that the pollen of *Poa annua* plants grown in cadmium-contaminated soil had an increased propensity to bind specific IgE (Aina et al., 2010). However, increased binding to IgE does not mean increased allergy symptoms, and only a certain proportion of patients sensitized will exhibit allergic symptoms (Galli et al., 2008). Further research is required to reveal the molecular cause and explanation for this observed phenomenon.

Therefore, we created a comprehensive approach for the comparison of pollen from polluted and environmentally preserved areas. To examine the effects of long-term, in vivo pollen exposure to multiple source pollutants, *Phleum pratense* (Timothy grass) pollen samples were collected along a regional road in Kruševac, central Serbia. This road experienced moderate traffic and was located near a chemical plant that produces fertilizers. Pollen samples from this location were compared




Fig. 1. Location of *Phleum pratense* pollen collection during the pollination seasons of 2014 and 2015 near Kruševac city, Serbia. Location P1 denotes a low population, rural environment that is free of industry and traffic pollution, whereas location P2 denotes a road traffic and industrially polluted environment.

with pollen samples collected from a rural, environmentally preserved area over two consecutive pollination seasons. We combined the quantitative comparison of proteome expression profiles from solution and 2D gels with unrestrictive in-depth quantitative PTM profiling using high resolution tandem mass spectrometry and the PEAKS 8.5 Suite platform. This was followed by quantitative IgE enzyme-linked immunosorbent assays (ELISA) and one dimensional (1D) IgE immunoblots that were probed with the sera of grass pollen allergic patients and healthy control subjects from Serbia. In addition, elemental and PAH compositional analyses of Timothy grass pollen samples from both locations, and the surface grain structure and SPP releasing potential (including total protein and phenolic content), were assessed.

2. Materials and methods

2.1. Pollen samples and collection areas

Timothy grass pollen samples were collected from two locations in central Serbia (Fig. 1) surrounding Kruševac (a city in central Serbia, with 75,000 inhabitants), with the shortest air-route direction being 10 km distance. One location was distant from the urban area in a non-polluted, environmentally preserved area (P1), with an air class of 1 or 2, i.e., good quality as per the U.S. Environmental Protection Agency (EPA) standard. The other location (P2) was beside the Kruševac-Kraljevo regional road and in close proximity to a chemical plant. Site P2 was thus exposed to both road traffic and chemical pollution, and had an average air quality of class 3 as per the EPA standard. The traffic road congestion score of 15% (which represents the increase in overall travel time relative to that of free flowing traffic) for this regional road is well below that of major European metropolises and big cities (https://www.tomtom.com/en_gb/trafficindex/). During 2011–2015, black carbon in the air was present on average for 5% of days per year (www.sepa.gov.rs). The average yearly concentrations of SO₂, NO₂, and O₃ for Kruševac city were 16.1 µg/m³, 24 µg/m³, and 12 µg/m³ (www.sepa.gov.rs), respectively, while the average yearly permissible limit was 50 µg/m³, 60 µg/m³ and 120 µg/m³, respectively (Matic et al., 2013). The highest daily concentrations of these gaseous pollutants during the same period were 10 times higher than the average yearly concentration during the 2012–2014 period (www.sepa.gov.rs).

Mature pollen grains, released by shaking from anthers, were collected in containers. The pollen released was sieved through stainless steel with a 325 mesh (44 µm) and stored at 4–8 °C prior to the extraction of pollen proteome. Nearly 25 g of Timothy grass pollen from each location was collected in each year during the sunny days within a seven-day frame of peak pollination season, with no or low wind speed below 1.5 m/s. The sampling was undertaken during the second week of July 2014 and the third week of July 2015, which are during the pollination summer seasons. The purity of the non-defatted *Phleum pratense* pollen samples from P1 and P2 were checked using particle counting, and were 99.0% and 98.5%, respectively. All procedures from the Timothy grass pollen collection onwards, including the purity check, were supervised and undertaken by the Institute for Virology, Vaccines and Sera, Torlak Institut, Belgrade, Serbia.

2.2. Analysis of released sub-pollen particles (SPPs)

The percentage of SPPs released at 5, 30, and 60 min from pollen grains deionized in water immersion in P1 and P2 samples from both seasons was determined using a Neubauer hemocytometer and light microscope (Motic, Kowloon, Hong Kong SAR) at 400× magnification that was connected to a Canon IXUS 400 digital camera for photo documentation with a further 5× magnification. The acquired digital images were analyzed with Adobe Photoshop, using the ruler function. A total of 350 ± 5 pollen grains from each *Phleum pratense* pollen sample were recorded (approximately 35–40 different microscopic fields/micrographs per pollen sample). This experiment was repeated

two times within a one-week time-frame for samples from both pollination seasons.

2.3. Pollen surface morphology assessment with scanning electron microscope (SEM)

After deposition of a thin gold layer onto the P1 and P2 Timothy grass pollen dust surface samples, a Mira 3 XMU field emission SEM (Tescan, Brno, Czech Republic) operated at 20 keV was used to analyze the surface morphology at 5000× to 50,000× magnification. The proportion of damaged grains was determined by visual counting. A grain was considered damaged if visible cracks of the exine were present or if it was broken (n = 300 ± 5 grains).

2.4. Preparation of Timothy grass aqueous pollen protein extract and determination of total protein content

Timothy grass pollen protein extracts were prepared as previously described (Ognjenovic et al., 2013), with minor modifications that reflect the standard procedures for the preparation of aqueous pollen extracts for diagnostic purposes (Vrtala et al., 1993). Briefly, P1 and P2 pollen samples (1 g each) were defatted with acetone, suspended in 10 mL of deionized water, and shaken overnight at 4 °C. The suspension was centrifuged at 14,000 ×g for 20 min at room temperature (RT) (20–25 °C). Protein sample concentrations were determined using the Bradford method (Bradford, 1976) and total protein amounts were assessed using 1D SDS-PAGE profiles and Image Quant TL version 8.1 software program (GE Healthcare, USA).

2.5. Measurement of total phenolic concentration

The method used to determine the concentration of ethanol-based phenolic extracts in P1 and P2 pollen samples was a modified version of the Folin-Ciocalteu protocol (Chun et al., 2003) that had been adjusted to allow for measurements in 96-well micro titer plates, as previously described by Mihajlovic et al. (2015). Total phenolics were expressed as milligrams of gallic acid equivalents (GAE) per gram of pollen.

2.6. Cohort of patients and ethics statement

Sera from 18 Serbian patients who were allergic to Timothy grass pollen, with variety of clinical respiratory symptoms such as allergic rhinitis (sneezing, runny/stuffy nose, and nasal congestion), allergic conjunctivitis (itchy, watery, and/or red eyes), and asthma (coughing, wheezing, chest tightness, and trouble breathing) were collected at the allergy clinic in the Institute for Virology, Vaccines and Sera, Torlak Institut, Belgrade, Serbia (Table S1, Supplementary information). Their specific IgE levels ranged from 12.2 to 126.0 kU_A/L; their average and median values were 36.1 and 17.1 kU_A/L, respectively (ImmunoCAP, g6; Phadia/Thermo Fisher, Uppsala, Sweden). The sera of two healthy individuals, one male and one female (< 0.1 kU_A/L), were used as the control. The study was approved by the National Ethics Committee of the University of Belgrade, Serbia (No. 017/6–990/66). Written informed consent was obtained from donors prior to blood donation and their data were processed and stored according to the principles stated in the Declaration of Helsinki. Sera were used individually for the 1D immunoblot test (first 16 patients), and all 18 patients were used individually for the quantitative ELISA test. The cohort of patients represented a 1:1 ratio of females to males, ranging in age from 25 to 51 years, with median and average ages of 34 and 36 years, respectively (Table S1).

2.7. Reagents and material

Organic solvents for mass spectrometry were purchased from J.T. Baker (Mallinckrodt Baker, Phillipsburg, NJ, USA). Ultra-pure water

(18 mΩ) was prepared using a Smart2Pure3™ Barnstead aqua purification system (Thermo Fisher Scientific, Waltham, MA, USA). Chemicals used for inductively coupled plasma optical emission spectrometry (ICP-OES) and for inductively coupled plasma mass spectrometry (ICP-MS) analysis were all analytical grade, and were supplied by Merck (Darmstadt, Germany). All other chemicals were purchased from Sigma-Aldrich (St Louis, MO, USA).

2.8. ICP-OES and ICP-MS measurements

Elements (excluding arsenic, mercury, indium, and cobalt) were measured using an ICP-OES 6500 Duo with a CID86 chip detector, controlled by the Iteva software program (Thermo Fisher Scientific, Waltham, MA, USA). Arsenic, mercury, indium, and cobalt were measured using an ICP-MS (ImmunoCAP Q, Thermo Scientific X series 2, Thermo Fisher Scientific, Waltham, MA, USA). The entire system was controlled by the Qtegra Instrument Control software program (Thermo Scientific, Waltham, MA, USA). Microwave digestion was performed using a microwave oven equipped with a rotor holding 10 PTFE cuvettes (Ethos 1, Advanced Microwave Digestion System, Milestone SRL, Sorisole, Bergamo, Italy). Two batches of polluted (P2) and environmentally preserved (P1) pollen from both pollination seasons were assessed.

2.9. Polycyclic aromatic hydrocarbon (PAH) determination

Pollen samples were prepared according to the Compendium Method TO-13A (U.S. Environmental Protection Agency (US EPA), 1999) using gas chromatography with a mass selective detector. PAHs were extracted in a microwave with a mixture of hexane and acetone (12.5 mL n-hexane:12.5 mL acetone) solvents, according to the EPA 3546 standard. The solvent volume was reduced to 1 mL by rotary evaporation under a reduced pressure atmosphere (55.6 kPa containing 0.2 mL isooctane as a keeper), then the n-hexane solution was reduced to 0.25 mL under a nitrogen stream at RT. All samples were analyzed with the internal standard method using a Agilent GC 6890 N (Agilent Technologies, Santa Clara, CA, USA), Agilent MSD 5973, and capillary column Restek (40 m × 0.18 mm × 0.07 μm; Restek, Bellefonte, PA, USA) in SIM mode.

Following PAH determination, compounds (external standards) were searched according to the US EPA set priority, as follows: naphthalene (Nap), acenaphthylene (Acy), acenaphthene (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr), benzo(a)anthracene (BaA), chrysene (Chry), benzo(b)fluoranthene (BbF), benzo(k)fluoranthene (BkF), benzo[a]pyrene (B[a]P), indeno(1.2.3-cd)pyrene (Ind), dibenzo(ah)anthracene (DahA), and benzo(ghi)perylene (BghiP). Ultra Scientific Semi-Volatiles Internal Standard Mixture ISM-560 with deuterated compounds, comprised of acenaphthene-d₁₀, chrysene-d₁₀, 1,4-dichlorobenzene, naphthalene-d₈, perylene-d₁₂, and phenanthrene-d₁₀, was used as an internal standard.

2.10. One dimensional (1D) and two dimensional (2D) SDS-PAGE

1D electrophoresis was performed on a 12% SDS polyacrylamide gel, following the standard Laemmli protocol (Laemmli, 1970) under reducing conditions. A total of 15 μg of *Phleum pratense* pollen proteins were loaded per well. A sample of 45 μg of pollen protein extracts were isoelectrofocussed on 7 cm IPG strips (GE Healthcare, Chicago, IL, USA), using an Ettan IPGphor 3 IEF System (GE Healthcare, Chicago, IL, USA) and further separated with 12% SDS-PAGE using a MiniProtean unit (Bio-Rad, Hercules, CA, USA). The 1D and 2D gels stained with colloidal Coomassie Brilliant Blue 250G were scanned with a Typhoon FLA 7000 (GE Healthcare, Chicago, IL, USA). The 2D gel spots were quantified and matched using the Image Master 2D Platinum v7.0 software program (GE Healthcare, USA).

2.11. Immunoblotting

The 1D acrylamide gels were transferred to PVDF membranes with a semi-dry Nova-Blot system (GE Healthcare, Chicago, IL, USA). The membranes were blocked in 1% bovine serum albumin (BSA) and 0.05% Tween 20 phosphate buffered saline (tPBS) for 2 h at RT. Sixteen patients shown in Table S1 were used for individual 1D blot analyses. Sera were diluted at a 1:10 ratio for 1D blot analysis in 0.2% BSA in tPBS. Membranes were incubated with sera at 4 °C, overnight with agitation, and then washed three times with tPBS. Immunoblot detection was performed with rabbit anti-human IgE antibody (dilution 1:2000; MIAB, Uppsala, Sweden) as secondary antibody (incubation at RT for 2 h), followed by an alkaline phosphatase conjugated goat anti-rabbit IgG (dilution 1:1000, incubation at RT for 2 h; Jackson ImmunoResearch, West Grove, PA, USA). The membrane strips were then simultaneously developed in 0.165 mg/mL 5-bromo-4-chloro-3'-indolylphosphate, 0.33 mg/mL nitro-blue tetrazolium in 100 mM NaHCO₃, and 5 mM MgCl₂, at a pH of 9.5. The 1D immunoreactive blots were scanned using Typhoon FLA 7000 (GE Healthcare, Chicago, IL, USA) and processed and normalized with the ImageQuant TL version 8 software program (GE Healthcare, Chicago, IL, USA).

2.12. In-gel and in-solution digestion procedures for mass spectrometry and shotgun proteomics analyses

After colloidal Coomassie Brilliant Blue staining and scanning, 2D gel spots were excised and in-gel digested with trypsin using the method described by Shevchenko et al. (Shevchenko et al., 2006). The proteins were digested with proteomics-grade porcine trypsin at a ratio of 1:20 (with between 25 and 75 ng of trypsin in 25 mM ammonium bicarbonate, depending on protein gel spot quantity). In-solution digestion of the short ragweed pollen fractions was performed according to the standard protocol, as previously described (Smiljanic et al., 2017).

2.13. Nano-LC-MS/MS

Trypsin-digested peptides were chromatographically separated using the EASY-nLC II system (Thermo Fisher Scientific, Waltham, MA, USA) with a 2-column set up: a trap column C18-A1, 2 cm (SC001, Thermo Fisher Scientific, Waltham, MA, USA) and analytical column PepMap C18, 15 cm × 75 μm, 3 μm particles, and 100 Å pore size (ES800, Thermo Fisher Scientific, Waltham, MA, USA). A total of 10 μL of each shotgun sample and 4 μL of 2D gel sample were loaded and separated, as previously described by Smiljanic et al., (2017).

2.14. Identification and label free quantification (LFQ) of Timothy grass pollen proteins and their post-translational modifications (PTMs)

Identification of Timothy grass pollen proteins and PTMs was performed using the PEAKS Studio 8.5 software program (Bioinformatics Solutions Inc., Waterloo, ON, Canada). Briefly, signature MS/MS spectra were searched using PEAKS DB and PEAKS PTM algorithms against a hybrid database consisting of a UniProtKB (tremble and reviewed) subfamily Pooideae database (taxon identifier 147,368, downloaded on August 17, 2017 from <http://www.uniprot.org/>) and cRAP (the common Repository of Adventitious Proteins) database (downloaded on October 18, 2016 from <http://www.thegpm.org/crap/>). Oxidation (Met) and deamidation (Gln, Asn) were considered as variables, with carbamidomethylation (Cys) set as fixed in the PEAKS DB algorithm. In the PEAKS PTM algorithm, an unrestricted PTM search was undertaken using an available list of 313 PTM items. Up to two missed trypsin cleavages with non-specific cleavages at both ends of a peptide were allowed. Mass tolerances were set to ± 10 ppm for parent ions and ± 0.5 Da for fragment ions. Protein filters were as follows: protein – 10 IgP ≥ 20, proteins unique peptides ≥ 1, and “A” Score for

confident PTMs identification of at least 20. Peptide filters were as follows: false discovery rate for peptide-spectrum matches < 0.5%; therefore, the resulting false discovery rate of the peptide sequence was lower than 1%, and de novo alignment local confidence score was ≥80%.

LFQ was performed for proteins and PTMs via PEAKS Q and PTM profiling algorithms (for PTMs quantification, normalization with protein level from LFQ data was performed) with default parameters and heat maps generated from P1 and P2 pollen samples. Each group consisted of three biological batches performed in technical duplicates, with P1 and P2 samples for season and extraction time matched.

2.15. Quantitative ELISA measurements

Quantitative IgE ELISA, data calculations, and conversion from absorbance at 450 nm to kU_A/L based on the interpolation of the linearly fitted standard curve with known amounts of standards in kU_A/L, were based on the methodologies described by Apostolovic et al. (2016) and Madhurantakam et al. (2010). Briefly, duplicates of seven

calibrators were prepared from an IgE myeloma with 640 kU_A/L of IgE (3-fold dilutions, with a range of 4.86–0.01 kU_A/L), and two control samples and patient samples were added to a 96-well plate and incubated for 2 h. The wells containing IgE calibrators and control samples were coated with 50 μL of 2.5 μg/mL monoclonal anti-human IgE (donated by Phadia, Uppsala, Sweden), and wells containing patient sera were coated with 50 μL of 5 μg/mL of the respective Timothy grass pollen protein extracts (P1 and P2) in 0.1 M carbonate buffer with a pH of 9.6. The secondary antibody, mouse anti-human IgE conjugated with horseradish peroxidase (Abcam, Cambridge, Cambs., UK), was incubated for 1 h. Finally, 3,3',5,5'-tetramethylbenzidine substrate was used to stop the reaction with 1 M H₂SO₄. The plates were read at 450 nm.

2.16. Statistical analyses

All experiments in this study were subjected to two-tail paired Student *t*-tests using GraphPad Prism 7.00 software (GraphPad, San Diego, CA, USA), to search for significant differences and descriptive




Fig. 2. A: Sub-pollen particle (SPP) release in *Phleum pratense* pollen samples from environmentally preserved (P1) and polluted (P2) locations measured at 5 min and 30 min time points. *** denotes significantly higher SPP release of P2 samples, compared with P1 samples, at *p* < 0.0001. B: Representative micrographs of both pollen samples using light microscopy during osmolysis in deionized water at 5 min, showing clear SPP release from P2 sample. Representative micrographs of P1 and P2 pollen samples in their dry, natural state, as well as ethanol-based extracts, are included to demonstrate the visual differences in the yellow color intensity and quality between them. C: SEM micrographs showing representative P1 and P2 pollen grains at a magnification of 45,000×. The red square in the micrograph of the P2 sample highlights the appearance of invaginations that are missing from the P1 sample. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

statistics. All analyses, except elemental content comparisons and patient sera immunoassays, contained at least 12 biological samples with technical duplicates for both P1 and P2 groups. Normality testing was performed using the built-in D'Agostino-Pearson normality test “omnibus K2” method in GraphPad Prism 7.00 program. Pearson correlation was applied to the IgE reactive responses of 18 grass pollen allergic patients run on P1 and P2 pollen samples obtained in the ELISA tests.

3. Results

3.1. Increased phenolic load and release of sub-pollen particles (SPPs) are prominent in pollen grains from polluted area (P2), compared with pollen grains from the preserved environment (P1)

A comparison of the ability of pollen samples to release SPPs is shown in Fig. 2A. After 5 min of pollen grain osmolysis in deionized water, 3% of pollen grains from P1 samples released SPPs, whereas P2 samples released almost 5 times more SPPs (14%) from pollen grains than the P1 samples (Fig. 2A). Thirty minutes after the aqueous incubation of pollen grains, P1 and P2 samples released SPPs from 17% and 37%, respectively, of the total number of pollen grains counted (Fig. 2A). Measurements from longer osmolysis times (e.g., 60 min) showed no further change in SPPs release from either sample (data not shown). Pollen samples and their pollen grains from both collection locations under light microscopy are shown in Fig. 2B, while representative differences in their grain surface as seen using scanning electron microscopy are shown in Fig. 2C.

Examination of the surface pollen grain morphology with SEM revealed that pollen in the environmentally preserved P1 samples had a regular, reticulate, net-like appearance with minute *spinuli*, whereas pollen grains from the polluted P2 samples contained frequent, irregular longer spines resembling *echinae* (Fig. 2C) that may represent areas where the exine wall is thinner.

The total protein content was not significantly different between P1 and P2 pollen samples (Fig. S1), whereas the content of phenolic compounds in ethanol extracts, as expressed in GAE, was significantly higher in P2 samples than in P1 samples (16.79 ± 0.05 mg GAE/g in P2 samples and 4.98 ± 0.06 mg GAE/g in P1 samples, at $p < 0.001$). Additionally, a more intense yellow color of ethanol extract was observed in P2 samples than in P1 samples (Fig. 2B), and the UV/Vis spectra had higher absorption maxima at 300–350 nm for the P2 samples (Fig. S2).

3.2. ICP-OES/MS measurements revealed significantly higher content of mercury, cadmium, and manganese in the pollen affected by air and road traffic pollution

The content of elements in the pollen samples are presented in Table 1.

The levels of heavy metals such as cadmium and mercury were significantly higher in the pollen samples from the polluted area (P2) than that from the environmentally pristine area (P1). In particular, there was a 25% higher content of cadmium in sample P2 than in sample P1, at $p = 0.0407$, and a 275% higher content of mercury in sample P2 than in sample P1, at $p = 0.0001$. Arsenic did not show excessive accumulation levels in either of the pollen samples, and the lead content also did not differ between pollen from the polluted P2 location and pollen from the environmentally pristine P1 location.

The levels of essential elements, such as iron, chromium, and copper, were significantly higher in pollen samples from the environmentally preserved location P1, and were 115% ($p = 0.005$), 70% ($p = 0.0014$), and 123% ($p < 0.0001$) higher, respectively, than that of the polluted location P2. In contrast, the manganese content of P2 samples were 54% higher than that of the P1 samples, at $p = 0.0025$. However, the overall results were comparable to the results obtained from unpolluted areas (Cloutier-Hurteau et al., 2014).

In contrast to these results, the PAH assessment did not reveal any air or road traffic pollution sources for pollen samples from either location (Table S2). From the 16 PAHs tested, only naphthalene and phenanthrene were detected, with exclusive presence of acenaphthylene and fluoranthene in P2 samples. However, these could be attributed to natural sources of pollution caused by wood or oil combustion (Mostert et al., 2010), because these are not markers for traffic pollution. Typical markers for traffic pollution instead would likely include benzo(ghi)perylene, benzo[a] pyrene, or similar higher molecular weight PAHs (Teixeira et al., 2012).

3.3. Label free quantification (LFQ) of Timothy grass pollen proteins from the environmentally preserved area and the chemical plant/road traffic polluted area

LFQ using the proprietary PEAKS software program enables quantitative comparison of two or more proteomes with their numerous peptides and protein constituents (Smiljanic et al., 2017). Three separate extractions were made of Timothy grass pollen samples from P1 and P2 during the 2014 and 2015 pollination seasons. Six different P1 samples matched to six P2 samples by season and time of extraction were run on an Orbitrap nLC-MS/MS system (Thermo Fisher Scientific, Waltham, MA, USA) as technical duplicates.

The most important and significant differences in protein abundances between the P1 and P2 pollen samples are shown in Fig. 3A. A comparison of the complete proteomes of P1 and P2 pollen samples as the percentage share of all allergen groups and all other groups is shown in Fig. 3B.

Oxidoreductase, superoxide dismutase (SOD), and peroxidase showed expression levels that were 2.7, 1.9, and 1.5 times higher, respectively, in P2 samples than that in P1 samples, at $p < 0.005$, $p < 0.01$, and $p < 0.05$, respectively. Glycogen phosphorylase was downregulated 2.6 times ($p < 0.005$), and the allergen group Phl p 12 was downregulated 2 times ($p < 0.01$), in P2 samples compared with P1 samples. In contrast, the allergen groups Phl p13 and Phl p 6 were downregulated 1.5 times ($p < 0.05$) in P2 samples compared with P1 samples (Fig. 3A).

The distribution or relative abundances of the allergen groups within each pollen sample (P1 and P2) is shown in Fig. 3B. This is

Table 1

Concentration of selected elements in the samples of *Phleum pratense* pollen determined by ICP-OES/MS.

Element ^a	Sample P1			Sample P2			p-Value
	Average	SD	N	Average	SD	N	
Li (µg/g)	0.13	0.06	8	0.05	0.01	8	0.0012
Mg (mg/g)	1.68	0.01	8	1.88	0.01	8	2.78E-20
Ca (µg/g)	1.24	0.34	8	1.25	0.17	8	0.9592
Sr (µg/g)	2.98	0.78	8	3.28	0.51	8	0.3701
Cr (µg/g)	1.65	0.34	8	0.97	0.34	8	0.0014
Mn (µg/g)	39.60	13.46	8	61.81	10.54	8	0.0025
Fe (µg/g)	155.78	67.52	8	72.79	20.83	8	0.0050
Ni (µg/g)	4.52	0.47	8	3.28	0.74	8	0.0531
Co (µg/g)	0.14	0.09	6	3.34	3.61	6	0.0557
Cu (µg/g)	16.18	3.93	8	7.24	0.60	8	1.78E-05
Zn (µg/g)	41.53	8.20	8	49.11	6.35	8	0.0578
As (µg/g)	0.04	0.17	5	0.04	0.03	7	0.8852
Cd (µg/g)	0.16	0.03	8	0.20	0.04	8	0.0407
In (µg/g)	0.64	0.07	5	0.56	0.18	5	0.3530
Hg (µg/g)	0.04	0.02	4	0.15	0.01	4	0.0001
Pb (µg/g)	0.15	0.04	7	0.19	0.08	7	0.2479

Bold font text denotes elements with significantly higher content in P2 samples than in P1 samples. N represents the number of samples analyzed from both pollination seasons.

^a Quantities are expressed as mg or µg of element per one gram of dry pollen material.




Fig. 3. Label free quantification (LFQ) of *Phleum pratense* pollen protein from environmentally preserved area (P1 samples) and chemical plant/road traffic polluted area (P2 samples), determined using PEAKS Suite 8.5. A: Protein groups with the highest and most significant differences in relative quantities between the P1 and P2 pollen samples. The areas under the extracted ion chromatogram (XIC) curve for each protein group in sample P1 are presented as 1, and P2 samples are expressed as P2/P1 XIC curve area ratios. B: Percentage share of allergen groups and redox enzymes of P1 and P2 pollen samples revealed by LFQ. All areas under the XIC curves of isoforms belonging to certain allergen group were summed and expressed as a percentage of the sum of total proteome of P1 or P2. * $p < 0.05$ and ** $p < 0.005$.

important because different allergens have different allergenic potencies. Some are considered to be major allergens (defined as an allergen that > 50% of allergic patients react with) and some are considered to be minor allergens; however, even the major allergens can have different IgE binding potency within a population. To date, there have been no quantitative proteomic studies, either absolute or relative, published for in-solution digested pollen protein extracts of Timothy grass.

3.4. Comparison of pollen proteomes of P1 and P2 reveals selective differences in certain protein abundancies in 2D SDS-PAGE maps, including qualitative difference in PTM profiling

Differences in the 2D electrophoretic profiles were pronounced, with a strikingly different pattern of appearances seen in the low molecular protein spots that are numbered in Fig. 4A. In addition, the overall appearance revealed a “smear” effect in the gel of the P2 samples, in contrast to the clearer and sharper protein spot contours found in P1 samples (Fig. 4A). Spots numbered in Fig. 4A were matched using the Image Master Platinum v 7.0 software program, and their prevalent proteins were identified (Table S3) as profilins (spots 1 and 5), major allergen Phl p 6 (spots 2 and 6; spots 3 and 7), and the major allergen grass group II/III Phl p 3 (spots 4 and 8). The larger area of matched spots that were darker in dye intensity in P2 samples was due to PTMs introduced by pollution and pollution-additive effects, not because of higher protein abundancies in the matched spots of the P2 sample gel, as revealed by the results from protein LFQ (Fig. 3A and B). Phl p 12, which belongs to the profilin group, was expressed almost two times as much in P1 samples compared with P2 samples (Fig. 3A); the allergen Phl p 6 group shows a similar pattern, but to a lesser extent. Representative examples of the most prevalent protein from matched

spots 1 (P1 sample) and 5 (P2 sample) are shown in Fig. 4B, with easily observable qualitative differences in the confident oxidative PTMs presence. Other than the PTMs identified in profilin 6 (pan-allergen Phl p 12, accession no. A4KA33) in the P1 samples, the PTMs found exclusively in P2 samples for the same isoform species were chlorination, carbamylation, and hexose binding (glycation or glycosylation, it remains to be elucidated precisely which), all on tyrosine residues and double oxidation of methionine (sulphone) (Fig. 4B).

3.5. Quantitative, in-depth profiling of PTMs in Timothy grass pollen proteomes

Applying the same setup used as for LFQ of pollen proteome extracts digested in solution, unrestricted searches and LFQ of PTMs were undertaken that included the labelled spots within 2D gels with a PEAKS PTM 8.5 algorithm. The oxidative PTMs are shown in Fig. 5 and Table S4, with relevant statistics for several PTM quantification parameters, such as the XIC curve area of modified and total amino acid residues, the percentage of modified residues out of the total, and the ratio obtained by dividing XIC curve areas of either P1 or P2 samples with the average XIC curve areas of P1 or P2 samples. In Fig. 5, each bar of the “Proteomes in Solution” section represents a single oxidative modification on a specified amino acid residue or residues as the sum of the averages of all the confidently identified and quantified PTMs of a certain type (e.g., the sum of all methionine oxidations throughout the complete proteome constitutes one sample).

Carbonylation is an irreversible protein modification, and it has been used as a biological marker for oxidative stress due to its early formation. Protein carbonylation typically occurs in three ways:

- Firstly, by direct oxidation with reactive oxygen species (ROS) on




Fig. 5. The relative abundance of oxidative post-translational modifications (PTMs) in environmentally preserved pollen samples (P1) and traffic polluted pollen samples (P2). One letter symbols represent amino acid residues: M represents Met; H, His; P, Pro; Y, Tyr; K, Lys; I, Ile; W, Trp; and F, Phe. P2/P1 ratio: this was obtained by dividing the sum of the averaged areas under the XIC curves of PTM from the P2 sample with the P1 sample. Numbers with percentages on graphs represent the percentage of modified amino acid residues out of total residues (100%), where the amount or total residues are composed of modified and unmodified parts. * denotes significantly higher XIC curve areas of P2 pollen samples than P1 pollen samples at $p < 0.05$. # denotes significantly higher percentages of modified residues of P2 pollen samples than P1 pollen samples at $p < 0.05$. Major protein allergen groups affected by designated PTMs are shown below the arrows. For more details, see Table S4.

observed in the 2D gel spots numbered 1 to 8 in Fig. 4A (Table S4). PTM quantification of the 2D gel spots is included as a semi-quantitative result, and the PTMs that could not be confidently detected in pollen proteome samples that had been digested in solution are shown in Fig. 5, such as the double oxidation of Met (sulphone); the dihydroxy form of Trp, Phe, and Tyr; the oxidation of Tyr to 2-amino Tyr; and the chlorination of Tyr. The quantification of PTMs from 2D gel proteins spots is not recommended, because the starting protein abundance in the matched spots are not the same and consequently unequal fluctuations in the extraction of trypsin-digested peptide concentrates between the P1 and P2 samples could be introduced during the processing of 2D spots. This is easily confirmed by observing the higher total XIC

curve areas of P2 samples, when compared with P1 samples; this contrasts with proteomes digested in solution, where the XIC curve areas of the total PTMs were almost equal and had no significant differences (Fig. 5, Table S4). This was confirmed by the quantitative nature of the in-solution trypsin digestion approach. However, the same trend of increased oxidative PTMs in P2 pollen was observed in samples designated as “proteomes in solution” and 2D gel spots (e.g., the oxidation of Met was 2.3 and 122 times higher, respectively, in P2 samples than in P1 samples; Table S4).

In addition, a set of PTMs designated as “physiological/spontaneous” (e.g., methylation, amidation, deamidation, deamidation followed by methylation, acetylation, and formylation) are shown in Table

S5, these were analyzed in the same manner as the PTMs shown in Fig. 5. None of these showed significant differences between P1 and P2 samples.

3.6. 1D immunoblots and ELISA testing of grass allergic patients results

The profiles of individual 1D immunoblots from 16 allergic patients performed on P1 and P2 pollen samples were very similar (Fig. 6). However, closer examination of the results revealed a higher molecular mass band (approximately 105 kDa), which is labelled as band 1 on the

diagram and the lowest IgE reactive band (band 7), which was below 14 kDa, were more prominent in P2 samples than in P1 samples. In contrast, the IgE reactive bands 5 and 6 at approximately 22 kDa and 17 kDa, respectively, were more intense in P1 samples than in P2 samples. Detailed results were obtained using the Image Quant TL v 8.1 densitometry function, followed by statistical analyses (Fig. 6). Significant differences (at $p < 0.05$) between the mean volumes of P1 and P2 samples were discrete, and ranged from as low as 10% of the volume range for IgE binding band 4 (from 35 to 25 kDa) to 30% for IgE reactive band 1. When interpreting these results, band 7 (10–14 kDa) was




Fig. 6. Representative 1D immunoblots of *Phleum pratense* pollen samples from an environmentally preserved area (P1) and a polluted area (P2) probed with the sera of grass pollen allergic patients (Table S1). The descending slope pictured above the immunoblot strips denotes that patient immunoblots are sorted in order of decreasing ImmunoCAP binding values, where g6 stands for the ImmunoCAP code of Timothy grass pollen extract. Lanes under the P1 and P2 gel designation are, from left to right: molecular weight markers; gel lanes of pollen extracts prior to Western blot analysis and, gel lanes of pollen extracts after Western blot analysis. The IgE responses of patients were normalized based on the difference in content of electrophoretic profiles prior to and after the Western blot analysis for P1 and P2 gels. These differences were expressed in μg , calculated based on known molecular marker concentrations and their total volume profiles. Numbers from 1 to 7 on the right side represent a position where major allergens or groups of allergens are present in all 16 Western blot strips in both P1 and P2 resolved pollen protein samples.

composed of Phl p 12 (profilins), Phl p 3, and Phl p 6 allergen groups (Table S3). While Phl p 2 and Phl p 3 allergens had the same abundances in both pollen samples (Fig. 3), there was a statistically significantly higher expression of Phl p 6 and 12 in the P1 samples of approximately 35% and 60%, respectively, than found in the P2 samples. Therefore, there was a higher IgE reactivity volume of approximately 15% in the P2 samples than in the P1 samples, and that of the P2 samples increased when lower quantities of allergen groups elicit a stronger IgE response. One possible explanation for this could be the significant load of oxidative PTMs found in the Phl p 6 and Phl p 3 allergens from the P2 samples (Figs. 4B and 5, Table S6). IgE binding band 3 (Fig. 6) was positioned at approximately 55 kDa and comprised allergen groups Phl p 4 (berberin bridge enzyme) and Phl p13 (polygalacturonases). The former was 1.24 times more abundant in P2 samples than in P1 samples, whereas the latter was 1.5 less abundant in P2 samples; therefore, the results counteracted each other, resulting in almost zero net effect, with difference in the means of 0.32 being almost 20% of the total volume range. No significant differences were found in either allergen groups by inspecting their individual PTMs profiles (Table S6). An Ole e 1-like allergen, Phl p11, that was found in IgE binding band 5 was equally abundant in both P1 and P2 pollen, constituting 0.8% of the total proteome, and there were no significant differences in IgE reactivity between P1 and P2 samples. Further study is required to determine the protein identities of allergenic band 1 (106–110 kDa), 3, and 6 (16–17 kDa). Phl p 1 and Phl p 5 are found within the 25–35 kDa range; both of these groups occupy almost equal portions of the P1 and P2 proteomes (Fig. 3). Again, IgE reactivity was higher in P2 samples, with the size of the significant mean difference comprising 10% of the volume range (Fig. 6). However, no significant differences in the specific PTMs profiling were found in either allergen group (Table S6).

Quantitative ELISA tests on protein extracts of P1 and P2 samples enabled IgE binding assessment of individual patient sera to pollen aqueous extracts under non-denaturing conditions (Fig. 7). In contrast to immunoblots that had subtle and differential responses for particular IgE binding bands and their continuous IgE epitopes, in the ELISA test all patients, with the exception of patients numbered 10 and 12, and showed stronger IgE binding in P1 samples than that in P2 samples. These ranged from 13% of the corresponding sample P2 kU_A/L value, to as much as 2.3 times the corresponding sample P2 kU_A/L value (Table S7). Both patient subgroups (high and moderate to low ImmunoCAP classes) exhibited significantly higher IgE binding values for P1 samples than for P2 samples (Fig. 7), which is almost the opposite trend to the results obtained from the 1D immunoblot (Fig. 6). This could be the consequence of the presence of heavy metal impurities or of increased polyphenol concentration in the P2 samples. In the case of immunoblots, these impurities are cleared during the electrophoretic resolution step. In summary, ELISA testing demonstrated superior or stronger IgE binding of patient sera to P1 pollen samples, in contrast to the variable and subtle changes observed in IgE reactivity from 1D immunoblots.

4. Discussion

In the present study, the effects of real, multiple source pollution patterns on *Phleum pratense* pollen were assessed by a comparison of the structure and releasing potential of SPPs; measurements of the levels of heavy metals, PAHs, and total polyphenolics in pollen; and quantitative profiling of proteome expression and PTMs, using pollen samples from the environmentally preserved area (P1). Additionally, the allergenic potential of the pollen was studied via IgE reactivity.

The bioavailability of pollen allergens is influenced by environmental factors such as light intensity, temperature, and humidity (Behrendt and Becker, 2001; Bryce et al., 2010). To minimize these confounding factors, pollen samples from both locations (P1 and P2) in the municipality of Kruševac, Serbia (Fig. 1), were collected at the same

time, and within a day of each other. Location P1 was positioned within a rural, less populated area (between Tekije and Dobromir) and was approximately 10 km away in a direct line from location P2. Location P2 was situated immediately along the Kruševac-Kraljevo regional road, and in the vicinity of a chemical plant that produces fertilizers. One of the main limitations of our study is the absence of sufficient evidence to support our assumption that location P2 is polluted and location P1 is preserved, since we were not able to conduct a comparative soil assessment, unlike in previous studies (Babic et al., 2009).

Pollen is exposed to contamination from substances and compounds due to the absorption of pollutants by plants from the air and soil, via the root systems for water-soluble pollutants, and by direct sedimentation of pollution from different origins. Therefore, pollen can be utilized as a bio-indicator of urban air pollution (Kalbande et al., 2008). Documenting heavy metal concentrations in pollen, and its effect on plant reproduction, may be essential to aid our understanding of how plants incorporate metals from metal-polluted sites.

The results of our analysis of the heavy metal content in pollen were in accordance with what has previously been reported in the literature (Cloutier-Hurteau et al., 2014; Xun et al., 2017). Many authors have emphasized that the level of cadmium in pollen from air polluted areas greatly exceeded the acceptable norms (Cloutier-Hurteau et al., 2014; Roman, 2009; Senechal et al., 2015). In our study, the cadmium concentration values of both P1 and P2 samples were several times higher than the highest acceptable concentrations for cadmium in pollen loads of 0.05 $\mu\text{g/g}$ intended for human consumption (Roman, 2009). Cadmium was 25% higher in P2 pollen samples than that in P1 pollen




Fig. 7. IgE binding of 18 patients allergic to Timothy grass pollen with high and low IgE antibody classes for *Phleum pratense* pollen extract (ImmunoCAP code: g6) as assessed by quantitative ELISA. A: Distribution of IgE binding values and their mean differences (represented by thick horizontal lines) from subjects in the high ImmunoCAP classes (left graph) and the moderate to low ImmunoCAP classes (right graph) to the environmentally preserved pollen (P1) and the chemical plant/road traffic polluted pollen (P2). B: Pearson correlation of quantitative ELISA assay results; r represents the Pearson correlation coefficient.

samples. However, the greatest significant difference in heavy metal content between P1 and P2 pollen samples was for mercury, which was 3.75 times higher in P2 samples and 4.5 times above the highest admissible content of heavy metals in pollen loads intended for human consumption (0.033 mg/kg) (Roman, 2009). Lead content was 20% higher in P2 samples than in P1 samples, although this difference was not statistically significant. However, lead was below the highest admissible concentration for lead in pollen intended for human consumption in both P1 and P2 samples (0.5 mg/kg); this could be due to the use of lead-free fuels in the last decade, compared with previous studies that have reported values above this concentration (Roman, 2009). Manganese had a 33% higher content in P2 samples than in P1 samples; however, no limits were determined on its extent in pollen loads. In contrast, the arsenic content was the same in both samples, which were both 5 times below the highest admissible concentration for arsenic in pollen intended for human consumption (0.2 mg/kg) (Roman, 2009).

The PAHs analyses did not reveal the presence of any traffic pollution markers, such as benzo (ghi) perylene, benzo [a] pyrene, or similar higher molecular weight PAHs. The only differentiating factors observed were acenaphthylene and fluoranthene. These are PAHs that are commonly formed during wood or oil combustion (Lambert et al., 2012).

Amino tyrosine, perhaps as a consequence of nitration and subsequent reduction by dithiothreitol treatment (Söderling et al., 2007), only occurred in tyrosine residue 72 in a few profilin isoforms, as determined using PTM profiling of 2D protein spots. These were more pronounced in P2 samples than in P1 samples. The nitration of tyrosine has been extensively studied, and is considered to be correlated with traffic pollution. This causes a wide range of effects and results in increased pollen allergenicity (Ackaert et al., 2014; Gruijthuijsen et al., 2006; Karle et al., 2012), yet in our study it was a rare event (0.6% of Tyr 72 residue in P2 profilin isoforms modified vs 0.2% in P1). The oxidation of tyrosine to 2-amino tyrosine could not be detected using a quantitative in-solution pollen proteomes approach but was detected in a more concentrated semi-quantitative manner using protein spots from the 2D gels. In contrast, seven different oxidative PTMs, which included serine O-hexose binding, were significantly more abundant in P2 pollen samples than in P1 samples. This highlights the oxidative stress induced by excessive levels of heavy metals (Emamverdian et al., 2015; Gzyl et al., 2015; Lounifi et al., 2013), since biotic and abiotic stresses are known to induce ROS production in plants (Apel and Hirt, 2004; Lounifi et al., 2013).

Oxidative stress induces various PTMs, with 35 types of peptide and protein oxidation described in the literature (Madian and Regnier, 2010). Some are reversible in vivo via enzymatic and non-enzymatic catalysis, but some are irreversible and happen spontaneously. Excessive oxidative stress leaves a protein carbonylation fingerprint in biological systems, and the extent of protein carbonylation is directly correlated to the time of exposure and the amount of ROS generated under stress conditions (Nguyen and Donaldson, 2005). Thus, these PTMs appear to be useful indicators for the severity of oxidative stress in plants. There are several challenges in detecting carbonylation PTMs, including their diversity, low presence, and instability (they readily react with Lys residues to form Schiff bases). This has led to several strategies for their identification and quantification, mainly via 2D digital gel electrophoresis in combination with mass spectrometry, preceded by either an enrichment and/or derivative strategy. However, besides these current limitations, these techniques are time and labor intensive, as well as expensive (Madian and Regnier, 2010).

While oxidative modifications such as the oxidation of Met (sulfoxide), and its double oxidation (sulphone), are spontaneous and reversible, they could be used to assess the effects of the exposure of a proteome to the influence of differential oxidative stress (e.g., different species and doses of ROS), as proposed by Holmstrom and Finkel (2014). In addition, the oxidation of histidine and the dihydroxylation

of phenylalanine and tryptophan may be useful as bio oxidative markers of intensive ROS action. In the present study, the oxidation of Met and His were 2.3 and 8.7 times higher, respectively, in polluted pollen (P2) than in environmentally preserved pollen (P1).

In the present study, we have shown a simple method for deep and wide PTM searches, which relies on in-solution trypsin-digested pollen proteomes and high resolution shotgun proteomic analyses with relative LFQ of the unrestricted PTM search. We managed to identify and confidently quantify over 25 different PTMs, with 11 being oxidative PTMs. Four of these were due to carbonylation, such as the oxidation of Lys to amino adipic semialdehyde, the oxidation or hydroxylation of Pro (direct way), and the Michael addition reaction of unsaturated aldehydes derived from lipid peroxidation as 4-hydroxy-2-nonenal (indirect way), which were all detected in abundant Phl p 6 allergens, with 6.0, 3.7, and 6.6 higher content, respectively, in P2 samples than in P1 samples (Fig. 5). These modifications have been associated with aging protein aggregation (Tanase et al., 2016). The glycosylation or glycation of tyrosine and serine residues with hexose (the identification of particular sugar involved still requires further research) as an indirect method for protein carbonylation was observed in Phl p 6 and profilins (Phl p 12 allergens), with 8 times higher content and an almost 10 times higher proportion of modified residues in P2 pollen samples than in P1 pollen samples. Further validation of our PTM search method is required. However, many of the criteria for the standardization of oxidative PTM quantification procedures expressed in the review paper by Nedic et al. (Nedic et al., 2015) have already been met and solved in our approach, which used relative LFQ and the advanced, proprietary PEAKS Suite 8.5 software program.

Carbamylation is a non-enzymatic and irreversible PTM that mainly results from the interaction between isocyanic acid and protein amino groups. Isocyanic acid is generally produced from the spontaneous decomposition of urea into ammonium and cyanate, a reactive species that is rapidly converted to isocyanic acid. We found that this PTM was 14.7 times more abundant in P2 pollen than in P1 pollen, and it accounted for 3.2% of all carbamylated residues (Lys, Asn, and Tyr), compared with 0.2% in P1 pollen samples (Table S6). The significance and physiological relevance of this finding, as well as that of levglandinyl-lysine anhydropyrrole adduct PTM (Table S5) and increased O-hexose binding in polluted pollen, remains unknown.

It is suggested that the Timothy grass from location P2 was subjected to with higher oxidative stress levels than that in location P1, caused by the higher concentrations of mercury and cadmium. This supposition was supported by evidence of the ROS chemistry scenario provided by the PTM profiling of the P1 and P2 pollen samples, and by a comparison of the proteome expressions between the two locations. The antioxidative defense enzymes were significantly upregulated, and it is reasonable to expect that SOD, oxidoreductase, and peroxidase would also be upregulated in response to chronic and increased oxidative stress caused by ROS.

In addition, total phenolics showed substantially higher (in excess of 3 times) extraction levels in P2 pollen samples than in P1 samples, again emphasizing the consequences of increased oxidative stress caused by mercury and cadmium contamination. This is because plants develop defense mechanisms in response to mercury and cadmium uptake by increasing the production of phenolic compounds, due to their radical scavenging properties and their ability to donate electrons (Manquian-Cerda et al., 2016). Total proteins decreased slightly in P2 pollen samples, though this change was not statistically significant; this is in agreement with previous studies on Timothy grass pollen pollution (Motta et al., 2006; Rogerieux et al., 2007). However, very different results have been obtained in other situations, which can be attributed to different plant species and different types and durations of pollution sources (Senechal et al., 2015).

We aimed to comprehensively study the changes in the Timothy grass pollen caused by air pollution that affect its allergenic potential, and these changes include not only the co- and PTMs of allergens and

proteins, but also chemical modifications as a direct consequence of reactions with pollutants, the quantity and expression profile of its proteome, and the metabolome. Important modifications to proteins are difficult to study in situ. Changes to the conformation of the proteins can occur as a consequence of extensive modifications, as can local denaturation of segments in the protein due to disruption of bonds that otherwise stabilize the overall structure.

When discussing the altered allergenicity of pollen grains, it is known that gaseous, traffic-related pollutants such as NO₂ (50 ppm, 4 h) and O₃ (0.7 ppm, 4 h) (Motta et al., 2006), as well as cadmium-contaminated soils (50 ppm, plant life time) (Aina et al., 2010), facilitate the release of allergen-rich SPPs from Timothy grass and *Poa annua* pollen. This subsequently increases the quantity of allergens in the respirable submicronic fraction (Aina et al., 2010; Motta et al., 2006). Results from the present study regarding the potential of plants to release SPPs are in accordance with previous studies (Motta et al., 2006; Aina et al., 2010), and showed that the capacity of polluted pollen to release SPPs under osmolytic conditions mimicking humid conditions and rain had more than doubled. This is likely due to the subtle changes seen in the exine that enabled them to fracture more easily, and to readily burst in water.

Finally, when comparing the IgE binding properties of P1 and P2 samples with Timothy grass allergic patients using quantitative ELISA tests, the P1 pollen samples were more potent. The sera from all patients, except for two out of the eighteen that reacted with the same intensity, showed at least a 15% increase in IgE binding in kU_A/L, and patients with ImmunoCAP classes 1–3 showed an even higher IgE binding increase (Table S7). Individual immunoblot results were divergent, showing a 10%–30% increase in IgE binding intensity of continuous epitopes with allergenic bands at approximately 105 kDa, 55 kDa (Phl p 4 and Phl p 13), 35 kDa & 25 kDa (Phl p 1 and Phl p 5), and 14 kDa & 10 kDa (Phl p 2/3, Phl p 6, and Phl p 12). The other three IgE binding band regions showed higher IgE binding potential in P1 pollen samples.

One possible explanation for the obtained disparate results between immunoblots and ELISA techniques is that discontinuous epitopes are more affected by the modifications than continuous epitopes, due to partial unfolding of important allergens. Oxidative conditions have been regarded as a cause of increased protein susceptibility to deamidation through increased flexibility of the polypeptide backbone or a transient unfolding of proteins, allowing asparagine or aspartic acid residues deamidation to occur and enhancing the formation of L-iso aspartyl residues (Cimmino et al., 2008). In addition, UV irradiation and heavy metals such as mercury and cadmium can cause an increase in the formation of ROS that leads to increased protein deamidation in cultured melanoma cells (D'Angelo et al., 2001), and to amyloid formation (Sadakane and Kawahara, 2018). These L-iso aspartyl residues are abnormal and can alter protein structure and function, as has been already shown for many proteins (Cimmino et al., 2008). To repair and restore these residues, cells use protein isoaspartyl carboxyl O-methyl transferases that can methyl esterify the free α-carboxyl groups of the isoaspartyl residues. Therefore, protein methylation following deamidation can be regarded as a marker of aspartate damage caused by excessive oxidative stress (Ingrosso et al., 2002). Our results are in agreement with those of previously published reports and show that deamidation was 2.2 times higher in P2 pollen than in P1 pollen, with significantly higher proportions of modified residues. Only rare, spontaneous, physiological PTMs with significant differences between P1 and P2 pollen samples were observed in our study, including deamidation followed by methylation modification, which was 2.5 higher in P2 pollen samples than in P1 samples (Table S5). In addition, polyphenols present in pollen can bind to allergens and disrupt important discontinuous IgE epitopes, making them hypoallergenic, as in the case of pomegranate juice and cashew nut allergens (Li and Mattison, 2018). Mercury is unusual among metals because it is capable of forming bonds with sulfur in cysteines that are similar to covalent bonds,

meaning that they are vulnerable protein targets. There is experimental evidence from Polacco et al. (2011) that has confirmed this direct effect of protein conformational disruption, by binding to exposed cysteine residues among mercury pleiotropic toxicity (Polacco et al., 2011).

Although the ELISA results showed a decrease in the IgE binding with polluted pollen, the increased content of oxidoreductases may increase the potential for polluted pollen to induce allergic reactions (Bacsi et al., 2006; Smiljanic et al., 2017). Additionally, there is increased potential for SPPs release by polluted pollen, which can further contribute to its allergenicity.

Heavy metal induced ROS species possess adjuvant effects in allergic diseases such as rhinitis and allergic asthma, including the increased capability for allergenic material release. The inhalation of polluted pollen that has increased heavy metal content is a threat for all individuals, not only those who are allergic (Madrigal et al., 2018). “*Sola dosis facit venenum*” (“The dose makes the poison,” a saying credited to Paracelsus), however, it is nearly impossible to deduce the toxic dose for healthy human nasal mucosa, lungs, or skin (Bolan et al., 2017). However, the cumulative effect of the increased levels of several heavy metals in polluted P2 pollen dramatically changed the amount of oxidative damage due to increased oxidative PTMs and deamidation, deamidation followed by methylation, as well as aberrant protein expression and derangements at cell physiology. Heavy metals, such as mercury, cadmium, and lead, have no physiological role in the human body. The increase in oxidative modifications, including deamidation, caused by heavy metal induced excessive ROS production, can either prove beneficial or lead to chronic diseases, autoimmune diseases (Mousavi et al., 2018), cancers (Chen et al., 2018), and accelerated aging in humans.

In conclusion, we have demonstrated that the presence of heavy metals and the consequent increased concentrations of phenolics in polluted grass pollen contributes to allergen modifications at the post-translational level. The increased presence of heavy metals leads to increased oxidative stress, which probably causes the excessive oxidative PTMs observed in this study; together with the increased concentrations of phenolics, this could disrupt important discontinuous epitopes in polluted pollen, and contribute to protein unfolding due to deamidation. This may be associated with higher rate of sensitization, with allergic responses due to changes in allergen structures, or due to its effect as an adjuvant. All Timothy grass pollen allergens are prone to oxidative modifications, with Phl p 6 emerging as the subject of multiple different oxidative modifications at several amino acid side residues. Further study is required to understand effects of polluted pollen on human health and allergy; hence, animal models of allergic diseases using this characterized pollen would be worthy of study.

Quantitative, unrestricted, and detailed PTM searches using an enrichment-free approach were used for the first time to map extensive modifications in the pollen allergome, which was shown to reflect the increased environmental oxidative stress. With some modifications, this PTM profiling approach is suitable for exploring the oxidative stress effects in any proteomic source in a quantitative in-depth manner, thus enabling further data-driven research.

Acknowledgements

This research work was funded the Ministry of Education, Science and Technological Development of the Republic of Serbia, GA No. OI172024, Belgian Special Research Fund BOF StG No. 01N01718, and the European Commission, under the Horizon 2020, FoodEnTwin project, GA No. 810752. The EC does not share responsibility for the content of the paper. This research was also supported by The Swedish Research Council, the Stockholm County Council (ALF project), the Swedish Asthma and Allergy Association's Research Foundation, the King Gustaf V's 80-year Foundation, the Swedish Heart-Lung Foundation, the Hesselman Foundation, the Konsul Th C Bergh Foundation, and the Swedish Cancer and Allergy Foundation.

Conflict of interest

Authors declare no conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2019.03.001>.

References

- Ackaert, C., Kofler, S., Horejs-Hoeck, J., Zulehner, N., Asam, C., von Grafenstein, S., Fuchs, J.E., Briza, P., Liedl, K.R., Bohle, B., Ferreira, F., Brandstetter, H., Oostingh, G.J., Duschl, A., 2014. The impact of nitration on the structure and immunogenicity of the major birch pollen allergen Bet v 1.0101. *PLoS One* 9, e104520.
- Aina, R., Asero, R., Ghiani, A., Marconi, G., Albertini, E., Citterio, S., 2010. Exposure to cadmium-contaminated soils increases allergenicity of *Poa annua* L. pollen. *Allergy* 65, 1313–1321.
- Apel, K., Hirt, H., 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399.
- Apostolovic, D., Sanchez-Vidaurre, S., Waden, K., Curin, M., Grundstrom, J., Gafvelin, G., Cirkovic Velickovic, T., Gronlund, H., Thomas, W.R., Valenta, R., Hamsten, C., van Hage, M., 2016. The cat lipocalin Fel d 7 and its cross-reactivity with the dog lipocalin Can f 1. *Allergy* 71 (10), 1490–1495.
- Babic, M., Filipovic-Trajkovic, R., Andjelkovic, S., Babic, S., 2009. The presence of heavy metals in soil near highway Krusevac—Kraljevo. *Protection of Nature* 60, 621–626.
- Bacsi, A., Choudhury, B.K., Dharajiya, N., Sur, S., Boldogh, I., 2006. Subpollen particles: carriers of allergenic proteins and oxidases. *J. Allergy Clin. Immunol.* 118, 844–850.
- Behrendt, H., Becker, W.M., 2001. Localization, release and bioavailability of pollen allergens: the influence of environmental factors. *Curr. Opin. Immunol.* 13, 709–715.
- Bolan, S., Kunhikrishnan, A., Seshadri, B., Choppala, G., Naidu, R., Bolan, N.S., Ok, Y.S., Zhang, M., Li, C.-G., Li, F., Noller, B., Kirkham, M.B., 2017. Sources, distribution, bioavailability, toxicity, and risk assessment of heavy metal(loid)s in complementary medicines. *Environ. Int.* 108, 103–118.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- Bryce, M., Drews, O., Schenk, M.F., Menzel, A., Estrella, N., Weichenmeier, I., Smulders, M.J., Buters, J., Ring, J., Gorg, A., Behrendt, H., Traidl-Hoffmann, C., 2010. Impact of urbanization on the proteome of birch pollen and its chemotactic activity on human granulocytes. *Int. Arch. Allergy Immunol.* 151, 46–55.
- Chen, W., Fu, W., Deng, Q., Li, Y., Wang, K., Bai, Y., Wu, X., Li, G., Wang, G., Huang, J., He, M., Zhang, X., Wu, T., Wei, S., Guo, H., 2018. Multiple metals exposure and chromosome damage: exploring the mediation effects of microRNAs and their potentials in lung carcinogenesis. *Environ. Int.* <https://doi.org/10.1016/j.envint.2018.11.020>.
- Chun, O.K., Kim, D.-O., Lee, C.Y., 2003. Superoxide radical scavenging activity of the major polyphenols in fresh plums. *J. Agric. Food Chem.* 51, 8067–8072.
- Cimmino, A., Capasso, R., Muller, F., Sambri, I., Masella, L., Raimo, M., De Bonis, M.L., D'Angelo, S., Zappia, V., Galletti, P., Ingrosso, D., 2008. Protein isospartate methyltransferase prevents apoptosis induced by oxidative stress in endothelial cells: role of Bcl-XL deamidation and methylation. *PLoS One* 3, e3258.
- Cloutier-Hurteau, B., Gauthier, S., Turmel, M.-C., Comtois, P., Courchesne, F., 2014. Trace elements in the pollen of *Ambrosia artemisiifolia*: what is the effect of soil concentrations? *Chemosphere* 95, 541–549.
- D'Angelo, S., Ingrosso, D., Perfetto, B., Baroni, A., Zappia, M., Lobianco, L.L., Tufano, M.A., Galletti, P., 2001. UVA irradiation induces L-isoaspartyl formation in melanoma cell proteins. *Free Radic. Biol. Med.* 31, 1–9.
- El Kelish, A., Zhao, F., Heller, W., Durner, J., Winkler, J.B., Behrendt, H., Traidl-Hoffmann, C., Horres, R., Pfeifer, M., Frank, U., Ernst, D., 2014. Ragweed (*Ambrosia artemisiifolia*) pollen allergenicity: SuperSAGE transcriptomic analysis upon elevated CO₂ and drought stress. *BMC Plant Biol.* 14, 176–192.
- Emamverdian, A., Ding, Y., Mokhberdorran, F., Xie, Y., 2015. Heavy metal stress and some mechanisms of plant defense response. *Sci. World J.* 2015, 756120.
- Galli, S.J., Tsai, M., Piliponsky, A.M., 2008. The development of allergic inflammation. *Nature* 454, 445–454.
- Grujthuisen, Y.K., Grieshuber, I., Stocklinger, A., Tischler, U., Fehrenbach, T., Weller, M.G., Vogel, L., Vieths, S., Poschl, U., Duschl, A., 2006. Nitration enhances the allergenic potential of proteins. *Int. Arch. Allergy Immunol.* 141, 265–275.
- Gzyl, J., Chmielowska-Bąk, J., Przymusiński, R., Gwóźdź, E.A., 2015. Cadmium affects microtubule organization and post-translational modifications of tubulin in seedlings of soybean (*Glycine max* L.). *Front. Plant Sci.* 6, 937.
- Holmstrom, K.M., Finkel, T., 2014. Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat. Rev. Mol. Cell Biol.* 15, 411–421.
- Ingrosso, D., Cimmino, A., D'Angelo, S., Alfinito, F., Zappia, V., Galletti, P., 2002. Protein methylation as a marker of aspartate damage in glucose-6-phosphate dehydrogenase-deficient erythrocytes: role of oxidative stress. *Eur. J. Biochem.* 269, 2032–2039.
- Ishizaki, T., Koizumi, K., Ikemori, R., Ishiyama, Y., Kushibiki, E., 1987. Studies of prevalence of Japanese cedar pollinosis among the residents in a densely cultivated area. *Ann. Allergy* 58, 265–270.
- Kalbande, D.M., Dhadse, S.N., Chaudhari, P.R., Wate, S.R., 2008. Biomonitoring of heavy metals by pollen in urban environment. *Environ. Monit. Assess.* 138, 233–238.
- Karle, A.C., Oostingh, G.J., Mutschlechner, S., Ferreira, F., Lackner, P., Bohle, B., Fischer, G.F., Vogt, A.B., Duschl, A., 2012. Nitration of the pollen allergen bet v 1.0101 enhances the presentation of bet v 1-derived peptides by HLA-DR on human dendritic cells. *PLoS One* 7, e31483.
- Kramer, U., Koch, T., Ranft, U., Ring, J., Behrendt, H., 2000. Traffic-related air pollution is associated with atopy in children living in urban areas. *Epidemiology* 11, 64–70.
- Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
- Lambert, O., Veyrand, B., Durand, S., Marchand, P., Le Bizec, B., Piroux, M., Puyo, S., Thorin, C., Delbac, F., Pouliquen, H., 2012. Polycyclic aromatic hydrocarbons: bees, honey and pollen as sentinels for environmental chemical contaminants. *Chemosphere* 86, 98–104.
- Li, Y., Mattison, C.P., 2018. Polyphenol-rich pomegranate juice reduces IgE binding to cashew nut allergens. *J. Sci. Food Agric.* 98, 1632–1638.
- Lounifi, I., Arc, E., Molassiotis, A., Job, D., Rajjou, L., Tanou, G., 2013. Interplay between protein carbonylation and nitrosylation in plants. *Proteomics* 13, 568–578.
- Madhurantakam, C., Nilsson, O.B., Uchtenhagen, H., Konradsen, J., Saarne, T., Høgbom, E., Sandalova, T., Gronlund, H., Achour, A., 2010. Crystal structure of the dog lipocalin allergen Can f 2: implications for cross-reactivity to the cat allergen Fel d 4. *J. Mol. Biol.* 401, 68–83.
- Madian, A.G., Regnier, F.E., 2010. Proteomic identification of carbonylated proteins and their oxidation sites. *J. Proteome Res.* 9, 3766–3780.
- Madrigal, J.M., Persky, V., Pappalardo, A., Argos, M., 2018. Association of heavy metals with measures of pulmonary function in children and youth: results from the National Health and Nutrition Examination Survey (NHANES). *Environ. Int.* 121, 871–878.
- Manquian-Cerda, K., Escudey, M., Zuñiga, G., Arancibia-Miranda, N., Molina, M., Cruces, E., 2016. Effect of cadmium on phenolic compounds, antioxidant enzyme activity and oxidative stress in blueberry (*Vaccinium corymbosum* L.) plantlets grown in vitro. *Ecotoxicol. Environ. Saf.* 133, 316–326.
- Matic, B., D., S., Jovanovic, D., Rakic, U., 2013. Report on Health Indicators of Environmental Protection of Republic of Serbia. 2012. pp. 1–60.
- Mihajlovic, L., Radosavljevic, J., Burazer, L., Smiljanic, K., Velickovic, T.C., 2015. Composition of polyphenol and polyamide compounds in common ragweed (*Ambrosia artemisiifolia* L.) pollen and sub-pollen particles. *Phytochemistry* 109, 125–132.
- Mostert, M.M.R., Ayoko, G.A., Kokot, S., 2010. Application of chemometrics to analysis of soil pollutants. *Trends Anal. Chem.* 29, 430–445.
- Motta, A.C., Marliere, M., Peltre, G., Sterenberg, P.A., Lacroix, G., 2006. Traffic-related air pollutants induce the release of allergen-containing cytoplasmic granules from grass pollen. *Int. Arch. Allergy Immunol.* 139, 294–298.
- Mousavi, S.E., Amini, H., Heydarpour, P., Amini Chermahini, F., Godderis, L., 2018. Air pollution, environmental chemicals, and smoking may trigger vitamin D deficiency: evidence and potential mechanisms. *Environ. Int.* <https://doi.org/10.1016/j.envint.2018.11.052>.
- Nedic, O., Rogowska-Wrzesinska, A., Rattan, S.I., 2015. Standardization and quality control in quantifying non-enzymatic oxidative protein modifications in relation to ageing and disease: why is it important and why is it hard? *Redox Biol.* 5, 91–100.
- Nguyen, A.T., Donaldson, R.P., 2005. Metal-catalyzed oxidation induces carbonylation of peroxisomal proteins and loss of enzymatic activities. *Arch. Biochem. Biophys.* 439, 25–31.
- Ognjenovic, J., Milcic-Matic, N., Smiljanic, K., Vuckovic, O., Burazer, L., Popovic, N., Stanic-Vucinic, D., Velickovic, T.C., 2013. Immunoproteomic characterization of *Ambrosia artemisiifolia* pollen allergens in canine atopic dermatitis. *Vet. Immunol. Immunopathol.* 155, 38–47.
- Parker, J.D., Akinbami, L.J., Woodruff, T.J., 2009. Air pollution and childhood respiratory allergies in the United States. *Environ. Health Perspect.* 117, 140–147.
- Polacco, B.J., Purvine, S.O., Zink, E.M., Lavoie, S.P., Lipton, M.S., Summers, A.O., Miller, S.M., 2011. Discovering mercury protein modifications in whole proteomes using natural isotope distributions observed in liquid chromatography-tandem mass spectrometry. *Mol. Cell. Proteomics* 10 (M110.004853-M004110.004853).
- Rogerieux, F., Godfrin, D., Senechal, H., Motta, A.C., Marliere, M., Peltre, G., Lacroix, G., 2007. Modifications of Phleum pratense grass pollen allergens following artificial exposure to gaseous air pollutants (O₃, NO₂, SO₂). *Int. Arch. Allergy Immunol.* 143, 127–134.
- Roman, A., 2009. Concentration of chosen trace elements of toxic properties in bee pollen loads. *Polish J. of Environ. Stud.* 18, 265–272.
- Ruokolainen, L., von Hertzen, L., Fyhrquist, N., Laatikainen, T., Lehtomaki, J., Auvinen, P., Karvonen, A.M., Hyvarinen, A., Tillmann, V., Niemela, O., Knip, M., Haahtela, T., Pekkanen, J., Hanski, I., 2015. Green areas around homes reduce atopic sensitization in children. *Allergy* 70, 195–202.
- Sadakane, Y., Kawahara, M., 2018. Implications of metal binding and asparagine deamidation for amyloid formation. *Int. J. Mol. Sci.* 19, 2449–2472.
- Schmidt, H., Gelhaus, C., Nebendahl, M., Janssen, O., Petersen, A., 2010. Characterization of Phleum pratense pollen extracts by 2-D DIGE and allergen immunoreactivity. *Proteomics* 10, 4352–4362.
- Senechal, H., Visez, N., Charpin, D., Shahali, Y., Peltre, G., Biolley, J.-P., Lhuissier, F., Couderc, R., Yamada, O., Malrat-Domenge, A., Pham-Thi, N., Poncet, P., Sutra, J.P., 2015. A review of the effects of major atmospheric pollutants on pollen grains, pollen content, and allergenicity. *Sci. World J.* 2015, 1–29.
- Shevchenko, A., Tomas, H., Havlis, J., Olsen, J.V., Mann, M., 2006. In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat. Protoc.* 1, 2856–2860.
- Smiljanic, K., Apostolovic, D., Trifunovic, S., Ognjenovic, J., Perusko, M., Mihajlovic, L., Burazer, L., van Hage, M., Cirkovic Velickovic, T., 2017. Subpollen particles are rich carriers of major short ragweed allergens and NADH dehydrogenases: quantitative proteomic and allergomic study. *Clin. Exp. Allergy* 47, 815–828.

- Söderling, A.-S., Hultman, L., Delbro, D., Højrup, P., Caidahl, K., 2007. Reduction of the nitro group during sample preparation may cause underestimation of the nitration level in 3-nitrotyrosine immunoblotting. *J. Chromatogr. B* 851, 277–286.
- Tanase, M., Urbanska, A.M., Zolla, V., Clement, C.C., Huang, L., Morozova, K., Follo, C., Goldberg, M., Roda, B., Reschiglian, P., Santambrogio, L., 2016. Role of carbonyl modifications on aging-associated protein aggregation. *Sci. Rep.* 6, 19311.
- Teixeira, E.C., Mattiuzi, C.D., Feltes, S., Wiegand, F., Santana, E.R., 2012. Estimated atmospheric emissions from biodiesel and characterization of pollutants in the metropolitan area of Porto Alegre-RS. *An. Acad. Bras. Cienc.* 84, 655–667.
- Vrtala, S., Grote, M., Duchene, M., van Ree, R., Kraft, D., Scheiner, O., Valenta, R., 1993. Properties of tree and grass pollen allergens: reinvestigation of the linkage between solubility and allergenicity. *Int. Arch. Allergy Immunol.* 102, 160–169.
- Xun, E., Zhang, Y., Zhao, J., Guo, J., 2017. Translocation of heavy metals from soils into floral organs and rewards of *Cucurbita pepo*: implications for plant reproductive fitness. *Ecotoxicol. Environ. Saf.* 145, 235–243.
- Zhao, F., Elkelish, A., Durner, J., Lindermayr, C., Winkler, J.B., Rusmali, R.F., Behrendt, H., Traidl-Hoffmann, C., Holzinger, A., Kofler, W., Braun, P., von Toerne, C., Hauck, S.M., Ernst, D., Frank, U., 2016. Common ragweed (*Ambrosia artemisiifolia* L.): allergenicity and molecular characterization of pollen after plant exposure to elevated NO₂. *Plant Cell Environ.* 39, 147–164.
- Zhao, F., Durner, J., Winkler, J.B., Traidl-Hoffmann, C., Strom, T.M., Ernst, D., Frank, U., 2017. Pollen of common ragweed (*Ambrosia artemisiifolia* L.): Illumina-based de novo sequencing and differential transcript expression upon elevated NO₂/O₃. *Environ. Pollut.* 224, 503–514.