

**Supplementary information for the article:**

Stojanović, M.; Lukić, I.; Marinković, E.; Kovačević, A.; Miljković, R.; Tobias, J.; Schabussova, I.; Zlatović, M.; Barisani-Asenbauer, T.; Wiedermann, U.; Inić-Kanada, A. Cross-Reactive Effects of Vaccines: Heterologous Immunity between Tetanus and Chlamydia. *Vaccines* **2020**, *8* (4), 719.

<https://doi.org/10.3390/vaccines8040719>.



**Supplementary Data : Table S1**

**Characteristics of selected anti-tetanus mAbs [35,36].**

mAb26	mAb51
<b>1. Affinity constant and rate of establishment the anti-TeNT mAb/TeNT interaction</b>	
$K_a = 0.4 \pm 0.2 \times 10^8 \text{ M}^{-1}$	$K_a = 2.4 \pm 0.4 \times 10^8 \text{ M}^{-1}$
$k_{on} = 1.1 \pm 0.2 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$	$k_{on} = 185 \pm 16 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$
Low affinity mAb	High affinity mAb
<b>2. Epitope positions on TeNT</b>	
Epitopes for mAb26 and mAb51 are located on close proximity of the ganglioside-binding place	
<b>3. Capability to inhibit TeNT/GD1b interactions</b>	
Limited inhibition of TeNT/GD1b interactions ( $29.9 \pm 3.1 \%$ )	Complete inhibition of TtNT/GD1b interactions ( $98.8 \pm 1.1 \%$ )
<b>4. In vivo protective potential against TeNT intoxication</b>	
Not protective against TeNT intoxication	Fully protective against TeNT intoxication